

- pyridine methiodide on mouse and rabbit hearts. *Acta Pharmacol Sin* 1989, 10 : 239
- 2 Bleeker WK, Mackaay AJC, Masson-Pévet M, Bouman LN, Becker AE. Functional and morphological organization of the rabbit sinus node. *Circ Res* 1980, 46 : 11
  - 3 Wang HZ, Wang DY, Zhou CM, Zhang KJ. Effect of nicotinamide on rabbit sinus node cell action potential and analysis of its mechanism. *Acta Pharm Sin* 1985, 20 : 340
  - 4 Wu RL, Wang YX, Zhu SM, Ge ZH. The recording of the transmembrane potential of the pacemaker cells in the isolated rabbit sinus node. *Acta Academiae Med Nanjing* 1984, 4 : 8
  - 5 Brown HF. Electrophysiology of the sinoatrial node. *Physiol Rev* 1982, 62 : 505
  - 6 Yanagihara K. Ionic current and pacemaker activity of the S-A node cells. *Jpn Circ J* 1980, 44 : 2618
  - 7 Nawrath H, Eick RET, McDonald TF, Trautwein W. On the mechanism underlying the action of D-600 on slow inward current and tension in mammalian myocardium. *Circ Res* 1977, 40 : 408
  - 8 Jin WZ, Zhou CM, Zhang KJ. Effects of nicotinamide on atrioventricular conduction on rabbits. *Acta Pharmacol Sin* 1984, 5 : 177
  - 9 Hatt H, Franke C. Taste receptors in crayfish: recording of single nicotinamide-activated channels. *Neurosci Lett* 1987, 73 : 137
  - 10 Goto J, Sperelakis N. Depression automaticity of the rabbit SA node by bepridil and nifedipine. *Eur J Pharmacol* 1984, 99 : 227
  - 11 Dangman KH, Hoffman BF. Effects of nifedipine on electrical activity of cardiac cells. *Am J Cardiol* 1980, 46 : 1059

\* \* \* \* \*

中国药理学报 *Acta Pharmacologica Sinica* 1989 Jul, 10 (4) : 339-342

## 二[二甲氨基]-二苯甲烷对豚鼠心乳头状肌动作电位的影响<sup>1</sup>

刘德强、赵德化、盛宝恒 (第四军医大学药理学教研室, 西安 710033, 中国)

### Effects of bis(dimethyl amino)-diphenyl methane (BDDM) on action potentials in guinea pig papillary muscle

LIU De-Qiang, ZHAO De-Hua, SHENG Bao-Heng

(Department of Pharmacology, The Fourth Military Medical College, Xi-an 710033, China)

**ABSTRACT** The effects of BDDM on action potentials and slow response action potentials in guinea pig papillary muscle were investigated by microelectrode technique. BDDM prolonged action potentials duration at 30, 50, 90, and 100% repolarization ( $APD_{30}$ ,  $APD_{50}$ ,  $APD_{90}$ ,  $APD_{100}$ ) and prolonged the effective refractory period (ERP), so ERP/

APD value became greater. BDDM (51.6  $\mu\text{mol/L}$ ) decreased the APA,  $V_{max}$  and prolonged  $APD_{50}$ ,  $APD_{90}$ . In barium-induced ventricular autorhythmicity, BDDM suppressed the maximal diastolic potential (MDP), APA and reduced the rate of spontaneous rhythm. These results suggest that BDDM may unspecifically inhibit the currents of  $\text{Ca}^{2+}$ ,  $\text{K}^{+}$ , and  $\text{Na}^{+}$ .

Received 1988 Jun 18 Accepted 1988 Nov 8  
<sup>1</sup> It was presented on the 5th Southeast and Western Pacific Regional Meeting of Pharmacologists, 1988 Jul, Beijing

**KEY WORDS** bis(dimethyl amino)-diphenyl methane (BDDM); papillary muscles; action potentials; barium

**摘要** 二[二甲氨基]-二苯甲烷(BDDM)能延长豚鼠心乳头状肌细胞APD、ERP,使ERP/APD比值增大;较高浓度时可降低APA和0相除极最大速率( $V_{max}$ );降低慢反应动作电位APA和 $V_{max}$ ,并延长APD<sub>90</sub>,抑制Ba<sup>2+</sup>诱发的乳头状肌自发电活动,降低APA和MDP。结果提示BDDM对心肌细胞Ca<sup>2+</sup>,K<sup>+</sup>,Na<sup>+</sup>的跨膜转运有抑制作用。

**关键词** 二[二甲氨基]-二苯甲烷;乳头状肌;动作电位;钡

4, 4'-二[二甲氨基]-二苯甲烷(4, 4'-bis(dimethyl amino)-diphenyl methane, BDDM)有抗实验性心律失常作用<sup>(1)</sup>,心室乳头状肌生理特性影响的实验提示<sup>(2)</sup>,此作用可能与其抑制钙内流有关。本文在离体豚鼠右心室乳头状肌上,用玻璃微电极技术,进一步观察BDDM对心室肌动作电位参数及不同慢反应电活动的影响,为BDDM的抗心律失常作用提供电生理学依据。

#### MATERIALS AND METHODS

豚鼠34只,♀♂不拘,体重328±SD 45g,击头致昏,速取心脏,放入经95%O<sub>2</sub>+5%CO<sub>2</sub>饱和的Tyrode液中,制备右心室乳头状肌标本,并将其以水平方向固定于2ml的浴槽内,用36±0.5℃,pH 7.2-7.4的Tyrode液循环灌注,速度5.5ml/min,以1Hz,3ms,1.5倍阈电压的刺激驱动标本。

记录电极为充以KCl 3mol/L的玻璃微电极,电阻10-40MΩ,微电极由Ag-AgCl丝与微电极放大器相连,输出信号分两路,一路输入双线示波器的上线进行观察,并输入光线示波器记录,另一路经电子微分器微分后输入双线示波器下线进行观察和光线示波器记录。标本稳定1h开始实验,一个标本只用一种药物,用南京无线电厂研制的博士-851型微电子计算机联机分析、处理图形,并打印结果。

BDDM白色粉末,兰州大学化学系陈淑英教授合成提供,实验前,先加少量稀盐酸溶解,再用蒸馏水稀释成1%的溶液,贮于4℃冰箱备

用。盐酸利多卡因(Lidocaine, Lid)天津人民制药厂。

#### RESULTS

**BDDM对豚鼠心室乳头状肌快反应动作电位参数的影响** BDDM 12.9 μmol/L作用于豚鼠心室乳头状肌标本20min后,动作电位时程(APD)缩短;BDDM 25.8, 51.6 μmol/L使APD<sub>100</sub>, APD<sub>90</sub>, APD<sub>50</sub>, APD<sub>30</sub>延长;BDDM 51.6 μmol/L还能使动作电位幅度(APA)及0相最大除极速率( $V_{max}$ )下降,膜电位(RP)负值减小(Tab 1)。

Tab 1. Effects of bis(dimethyl amino)-diphenyl methane (BDDM) on action potential parameters and effective refractory period (ERP) in guinea pig papillary muscles. n=6,  $\bar{x}$ ±SD. \*P>0.05, \*\*P<0.05, \*\*\*P<0.01.

Parameter	BDDM (μmol/L)			
	Control	12.9	25.8	51.6
APA(mV)	125±5	125±5*	123±6*	120±7***
RP(mV)	89±2	88±1*	88±2*	86±2*
$V_{max}$ (V/s)	271±41	270±37*	251±31*	234±43**
APD <sub>30</sub> (ms)	101±11	98±9**	109±7**	107±11**
APD <sub>50</sub> (ms)	148±11	144±13**	162±11***	167±11***
APD <sub>90</sub> (ms)	202±18	194±19***	217±19**	229±17***
APD <sub>100</sub> (ms)	224±17	217±19***	243±20***	254±18***
ERP(ms)	160±23	163±19*	183±26***	205±34***
ERP/APD	0.7±0.1	0.8±0.1***	0.8±0.1***	0.8±0.1***

**BDDM对有效不应期(ERP)的影响** 在本刺激节律为1Hz的基础上,每8次刺激插入1次期前收缩,逐步缩短期前刺激和基本节律刺激之间的间隔,以期前刺激能引起动作电位产生的最短时间间隔为ERP,以上方法由电子计算机控制输出。BDDM 12.9 μmol/L,可使豚鼠心乳头状肌ERP缩短,但ERP/APD比值增大;BDDM 25.8, 51.6 μmol/L使ERP/APD比值增大(Tab 1)。

**BDDM 对高 K<sup>+</sup> 除极慢反应动作电位的影响** 提高灌流液中 K<sup>+</sup> 浓度为 18 mmol/L, NaCl 相应降低, 以保持溶液渗透压不变。以刺激强度大于正常阈值 4 倍, 波宽 8 ms 的电刺激也不能使豚鼠心室乳头状肌产生动作电位, 证实此时快钠通道基本失活<sup>(3)</sup>, 给异丙肾上腺素 1 μmol/L, 改用 0.25 Hz, 5 ms, 2 倍阈电压刺激, 又出现动作电位, 其 V<sub>max</sub> 为 10.0 ± 1.9 V/s (n=7), 呈慢反应型<sup>(4)</sup>。BDDM 25.8, 103.2 μmol/L 可使慢反应动作电位之 APA 及 V<sub>max</sub> 降低, APD<sub>50</sub> 延长 (Tab 2)。

Tab 2. Effect of BDDM on slow response action potentials pretreated with KCl 18 mmol/L in guinea pig papillary muscles, n=7,  $\bar{x} \pm SD$ . \*P>0.05, \*\*P<0.05, \*\*\*P<0.01.

Parameter	BDDM (μmol/L)		
	Control	25.8	51.6
APA (mV)	72 ± 7	68 ± 6**	63 ± 9***
V <sub>max</sub> (V/s)	10 ± 2	9 ± 2***	8 ± 2***
APD <sub>50</sub> (ms)	69 ± 8	72 ± 9*	78 ± 13***
APD <sub>90</sub> (ms)	123 ± 15	132 ± 17**	141 ± 23***

#### BDDM 对 Ba<sup>2+</sup> 诱发异常自律性的影响

在灌流液中加入 BaCl<sub>2</sub> 3 mmol/L 时, APA 降低, 约 20 min 后出现自发活动, 舒张电位从 -80 mV 上移到 -50 mV 左右。BDDM 25.8 μmol/L 作用于豚鼠乳头状肌标本 (n=5) 20 min 后, APA 从 72 ± 4 降低到 66 ± 4 mV (P<0.01); 自动发放频率由 105 ± 9, 降低到 93 ± 9 bpm (P<0.01); BDDM 51.6 μmol/L 使 APA 由 72 ± 4 降到 60 ± 5 mV (P<0.01); MDP 从对照的 55 ± 4 降到 46 ± 9 mV (P<0.01), 频率从 105 ± 9 降到 86 ± 8 bpm (P<0.01)。Lid 55.4, 110.8 μmol/L 使自发频率从 98 ± 4, 分别降到 85 ± 6, 77 ± 5 bpm (P 均 < 0.01)。

#### DISCUSSION

药物对心肌细胞的电生理影响, 是抗心律失常药的作用基础<sup>(5)</sup>。BDDM 可使 APD, ERP 延长, ERP/APD 比值增大, 这些作用能使单

向传导阻滞变为双向传导阻滞, 有利于中止折返冲动<sup>(6)</sup>。V<sub>max</sub> 减低能阻滞冲动在未完全复极的心肌纤维内传导, 发挥其抗心律失常作用。较高浓度的 BDDM 能降低心肌细胞的 APA 和 V<sub>max</sub>, 说明 BDDM 对 I<sub>Na</sub> 有抑制作用。

内向整流电流 (I<sub>K</sub>) 的减少, 与慢内向电流 (I<sub>si</sub>) 的增加, 是形成心室肌动作电位坪台期的主要原因<sup>(7)</sup>。维拉帕米等慢通道阻滞剂, 可使动作电位时程延长, 其原因是复极化时, K<sup>+</sup> 外流可由于细胞内 Ca<sup>2+</sup> 减少而减少<sup>(8)</sup>, Hoffman 等也证明, 当细胞外 Ca<sup>2+</sup> 减少时, 动作电位时程延长<sup>(9)</sup>。BDDM 小剂量使动作电位时程缩短, 与阻滞心肌细胞的钙内流有关, BDDM 25.8, 51.6 μmol/L 也使动作电位时程延长; 在高 K<sup>+</sup> 除极后慢反应动作电位中, BDDM 能降低 APA 和 V<sub>max</sub>, 延长动作电位时程。提示 BDDM 不但能阻滞心肌细胞的 Ca<sup>2+</sup> 内流, 而且能降低心肌细胞的 K<sup>+</sup> 外流。

Ba<sup>2+</sup> 诱发的自发性电活动主要是通过降低钾电导, 增加跨膜净 I<sub>si</sub> 内流所致<sup>(10)</sup>。BDDM 和 Lid 均能使 Ba<sup>2+</sup> 诱发的慢反应型自发电活动的频率减慢, BDDM 还能使其 APA 及 MDP 降低, 也提示 BDDM 具有 Ca<sup>2+</sup> 通道阻滞作用。以上实验结果表明, BDDM 对心肌细胞的 Ca<sup>2+</sup>, K<sup>+</sup>, Na<sup>+</sup> 跨膜转运均表现抑制作用。

#### REFERENCES

- 1 Liu DQ, Zhao DH, Sheng BH. Antiarrhythmic effects of bis (dimethyl amino)-diphenyl methane (BDDM). *Acta Pharm Sin* 1988; 23 : 448
- 2 Liu DQ, Zhao DH, Sheng BH. Effects of bis (dimethyl amino)-diphenyl methane (BDDM) on physiological properties of isolated guinea pig papillary muscles. *Chin J Pharmacol Toxicol* 1989; 3 : 49
- 3 Fan Z, Liang ZJ, Yang XY. Effect of sodium tanshinone II-A sulfonate on electrical and mechanical properties of myocardium. *Acta Pharmacol Sin* 1986; 7 : 527
- 4 Noma A, Irisawa H. The effect of sodium ion on the initial phase of the sinoatrial pacemaker action potential in rabbits. *Jpn J*

- Physiol* 1974; 24 : 617
- 5 Hauswirth O, Singh BN. Ionic mechanisms in heart muscle in relation to the genesis and the pharmacological control of cardiac arrhythmias. *Pharmacol Rev* 1978; 30 : 5
  - 6 Manzanares J, Tamargo J. Electrophysiological effects of imipramine in non-treated and in imipramine-pretreated rat atrial fibres. *Br J Pharmacol* 1983; 79 : 167
  - 7 Goodman AG, Gilman LS. *The pharmacological basis of therapeutics*. 6th ed. NY: Macmillan, 1980: 762
  - 8 Isenberg G. Cardiac Purkinje fibres: the slow inward current component under the influence of modified  $[Ca^{2+}]_i$ . *Pflugers Arch* 1977; 371: 61
  - 9 Hoffman BF, Suckling EE. Effect of several cations on transmembrane potentials of cardiac muscle. *Am J Physiol* 1956; 186 : 317
  - 10 Hay DWP, Wadsworth RM. The effects of calcium channel inhibitors and other procedures affecting calcium translocation on drug-induced rhythmic contractions in the rat vas deferens. *Br J Pharmacol* 1983; 79 : 347

\* \* \* \* \*

中国药理学报 *Acta Pharmacologica Sinica* 1989 Jul; 10 (4) : 342-345

## 达唑氧苯对麻醉兔脑血管阻力的作用<sup>1</sup>

芮耀诚、孙笃新、龙 焜 (第二军医大学药理学系药理教研室, 上海 200433, 中国)

### Effect of dazoxiben on cerebrovascular resistance in rabbits

RUI Yao-Cheng, SUN Du-Xin, LONG Kun (*Department of Pharmacology, Faculty of Pharmacy, Second Military Medical University, Shanghai 200433, China*)

**ABSTRACT** The effects of dazoxiben, a  $TXA_2$  synthetase inhibitor, and indomethacin were compared on cerebrovascular resistance (CVR) and levels of serum  $TXB_2$ , 6-keto-PGF<sub>1 $\alpha$</sub>  (the stable metabolites of  $TXA_2$  and PGI<sub>2</sub>, respectively) and on protection from acute brain ischaemia caused by *ia* arachidonic acid (AA) in rabbits. The flow represented the cerebral blood flow (CBF) in two internal jugular arteries were measured with electromagnetic flow meter after occlusion of bilateral vertebral arteries and external jugular arteries. CVR was represented as blood pressure/(CBF·100 g brain). Serum  $TXB_2$  and 6-keto-PGF<sub>1 $\alpha$</sub>  levels were determined by radio-

immunoassay. The results showed that CVR and BP, EEG, ECG were not affected by treatment with *iv* dazoxiben 2 or 10 mg/kg. The CVR was enhanced by 35.5 and 49.8% at 30 and 40 min, respectively after *iv* indomethacin 10 mg/kg. The serum  $TXB_2$  level ( $872 \pm 85$ ) was inhibited to  $511 \pm 169$  pg/ml ( $n=5$ ,  $P<0.05$ ) and 6-keto-PGF<sub>1 $\alpha$</sub>  increased from  $668 \pm 309$  to  $890 \pm 357$  pg/ml ( $n=5$ ,  $P<0.05$ ) at 30 min after *iv* 2 mg/kg dazoxiben. However, both  $TXB_2$  and 6-keto-PGF<sub>1 $\alpha$</sub>  decreased by 26.4 and 32.7%, respectively at 40 min after *iv* indomethacin 10 mg/kg. In a model of cerebral ischaemia caused by *ia* AA in rabbits, the EEG change and enhancement of CVR were antagonized by *iv* dazoxiben 10 mg/kg completely, but only partly antagonized by indomethacin 10 mg/kg. The

Received 1988 Jul 23 Accepted 1988 Dec 28

<sup>1</sup>Project supported by the Shanghai Science and Technology Developing Foundation № 863430906