

Scand 1980; 62 : 345-56.

eg-Olofsson O. Hypoglycaemia and neurological disturbances in children with diabetes mellitus. *Acta Paediatr Scand* 1977; 270 (Suppl) : 91-5.

14 MacCuish AC, Munro JF, Duncan LJP. Treatment of hypoglycaemic coma with glucagon, intravenous dextrose and mannitol infusion in a hundred diabetics. *Lancet* 1970; 2 : 946-9.

LD: ISSN 0253-9756 中国药理学报 *Acta Pharmacologica Sinica* 1992 May; 13 (3) : 197-200

## Effects of Picroliv, the active principle of *Picrorhiza kurroa*, on chemical changes in rat liver poisoned by *Amanita phalloides*

Shreshth DWIVEDI<sup>1</sup>, Ravi RASTOGI, Narendra Kumar GARG, Bhola Nath DHAWAN<sup>2</sup>  
 MR Centre for Advanced Pharmacological Research on Traditional Remedies, Central Drug Research Institute, Lucknow - 226001, India)

**ABSTRACT** The efficacy of Picroliv, a standardized iridoid glycoside fraction of *Picrorhiza kurroa*, studied against the *Amanita phalloides*-induced chemical changes in rat liver. *A phalloides* (50 mg · kg<sup>-1</sup>) caused significant increases in the activities of hepatic 5'-nucleotidase,  $\gamma$ -glutamyl transaminase, acid ribonuclease, and succinate dehydrogenase, but a decrease in glucose-6-phosphatase. The level of cytochrome P-450 in microsomal fraction and content of glycogen in liver showed significant decreases. Picroliv (25 mg · kg<sup>-1</sup> · d<sup>-1</sup> × 10 d) provided significant restorations of all the biochemical changes induced by *A phalloides* except cytochrome P-450 and glycogen. These results demonstrated the protective effect of Picroliv against *A phalloides*-induced toxicity in rats.

**KEY WORDS** *Amanita*; *Picrorhiza kurroa*; Picroliv; liver; enzyme tests; cytochrome P-450; glycogen

Picroliv (Pic) is a standardized iridoid glycoside fraction isolated from roots and

rhizomes of *Picrorhiza kurroa*. It contains 60% picroside I and kutkoside in a ratio of 1 : 1.5, the balance being minor constituents. Earlier studies have shown the hepatoprotective efficacy of Pic against CCl<sub>4</sub><sup>(1)</sup>, galactosamine<sup>(2)</sup>, monocrotaline<sup>(3)</sup>, paracetamol<sup>(4)</sup>, and thioacetamide<sup>(5)</sup> induced liver damages in rats, and *Plasmodium berghei* infection in *Mastomys*<sup>(6)</sup>.

*A phalloides* (AP) is the most toxic mushroom causing 95% of total mushroom poisonings. It contains mainly 2 toxins : amanitin and phalloidin. Phalloidin is the major toxic agent and is known to act on plasma membrane of hepatocytes in rats<sup>(7)</sup>. However, the mortality of rats poisoned with AP was reduced by Pic<sup>(8)</sup>. In this communication we have reported the influence of Pic on biochemical changes in rats poisoned by AP in rat liver.

### MATERIALS AND METHODS

Adult ♂ rats (wt 130 ± s 15 g) of Sprague-Dawley strain, inbred in CDRI Animal House were used. The rats were fed *ad lib* standard pellet diet (Lipton, Bombay) and allowed free access to water.

Received 1991 Sep 11 Accepted 1992 Jan 17  
 Senior Research Fellow of the Council of Scientific and Industrial Research, New Delhi.  
 Correspondence: Prof B N Dhawan, FNA, Director, Central Drug Research Institute, Lucknow - 226001, India.

**Induction of hepatotoxicity** Rats were injected ip *AP* ( $50 \text{ mg} \cdot \text{kg}^{-1}$ ), fasted overnight and decapitated after 10 d.

**Treatment with Pic** A freshly prepared aqueous solution of Pic was given intragastrically (ig) at a dose of  $25 \text{ mg} \cdot \text{kg}^{-1}$  simultaneously with *AP* injection. Pic feeding was continued for 9 d further. Two groups of rats were used as control, one receiving normal saline with *AP* and the other an equal volume of normal saline only.

Before decapitation, blood was drawn from the retro-orbital plexus and kept for 30 min to obtain the serum. The liver was excised, washed with chilled  $\text{NaCl } 145 \text{ mmol} \cdot \text{L}^{-1}$  (solution), and 10% (wt/vol) homogenate was prepared in  $\text{KCl } 154 \text{ mmol} \cdot \text{L}^{-1}$ .

**Biochemical estimations** The activities of 5'-nucleotidase (EC 3.1.3.5),  $\gamma$ -glutamyl transpeptidase (EC 2.3.2.2), acid ribonuclease (EC 3.1.4.22), acid phosphatase (EC 3.1.3.2), succinic dehydrogenase (EC 1.3.99.1), glucose-6-phosphatase (EC 3.1.3.9), cytochrome P-450, and cytochrome  $b_5$  in liver and GOT, GPT, and alkaline phosphatase in serum were assayed as described in Ref 4.

The levels of lipid peroxides, total lipids, phospholipids, cholesterol, DNA, RNA, total proteins, and liver glycogen and total serum protein and albumin were also measured as in Ref 4.

**Statistical analyses** The *AP*-treated group was compared with the control while the Pic-treated group was compared with *AP*-intoxicated group. One way analysis of variance was done among 3 groups. Individual significance of group means was done by Newman's and Keul's test for multiple comparison. *P* values were calculated by using *t* test.  $P < 0.01$  was considered significant.

## RESULTS

**Induction of hepatotoxicity by *AP*** Pilot experiments were carried out at 2.5, 5.0 (iv)

25, 50 (ip)  $\text{mg} \cdot \text{kg}^{-1}$ . The changes in rat liver and serum were studied after 1, 2, 10, and 20 d. The magnitude of changes was maximum on d 10 after  $50 \text{ mg} \cdot \text{kg}^{-1}$ . The biochemical changes in liver poisoned by *AP* and their values are given in Tab 1.

*AP* caused significant augmentation in the activities of 5'-nucleotidase and  $\gamma$ -glutamyl transpeptidase. The increase in the activity of 5'-nucleotidase was relatively slight (18%) but that of  $\gamma$ -glutamyl transpeptidase was much higher (60%). About 50% elevation in the activities of acid ribonuclease and succinic dehydrogenase was observed. The activity of glucose-6-phosphatase in liver and content of cytochrome P-450 in the microsomal fraction of liver cells were decreased significantly, the degree of reduction being 43% and 78%, respectively. Other hepatic enzymes acid phosphatase [ $55 \pm 3 \text{ nmol of } p\text{-nitrophenol formed} / (\text{min} \cdot \text{mg protein})$ ] and cytochrome  $b_5$  [ $0.31 \pm 0.02 \text{ nmol} / (\text{mg microsomal protein})$ ] remained unaffected. The hepatic glycogen was found to be almost completely exhausted. However, other chemical constituents of liver like total lipids ( $19.5 \pm 4 \text{ mg} \cdot \text{g}^{-1}$ ), phospholipids ( $19.5 \pm 0.4 \text{ mg} \cdot \text{g}^{-1}$ ), cholesterol ( $9.7 \pm 0.3 \text{ mg} \cdot \text{g}^{-1}$ ), lipid peroxides ( $117.5 \pm 26.0 \text{ nmol of malonaldehyde formed} \cdot \text{g}^{-1}$ ), DNA ( $2.6 \pm 0.1 \text{ mg} \cdot \text{g}^{-1}$ ), RNA ( $4.38 \pm 0.37 \text{ mg} \cdot \text{g}^{-1}$ ), and proteins ( $164 \pm 5 \text{ mg} \cdot \text{g}^{-1}$ ) remained unaffected. Serum parameters did not show any significant change following *AP* exposure. The levels of cytochrome  $b_5$ , DNA, RNA, protein, total lipids, phospholipids, cholesterol, and lipid peroxides in liver and GOT, GPT, alkaline phosphatase, bilirubin, protein, and albumin in serum were not altered at any of the doses of *AP* during 1-20 d.

**Hepatoprotection by Pic** Pic  $25 \text{ mg} \cdot \text{kg}^{-1}$  for 10 d, significantly reversed the increase in the activities of 5'-nucleotidase

Tab 1. Effects of Picroliv on biochemical indices in liver poisoned by *Amanita phalloides* and Q values with significances.  $n=6$  rats.  $\bar{x} \pm s$ . \*\* $P < 0.01$  vs Control. \* $P > 0.05$ , \*\*\* $P < 0.01$  vs *A phalloides*. §  $P > 0.05$ , §§§ $P < 0.01$  vs *A phalloides* + Picroliv. (% of protection in parentheses)

	Control	<i>A phalloides</i>	<i>A phalloides</i> + Picroliv
5'-Nucleotidase, nmol P <sub>i</sub> released / (min · mg protein)	38 ± 2 Q = 0.94 §	45 ± 1 Q = 11.79***	39 ± 1 (86%) Q = 10.85***
γ-Glutamyl transpeptidase, ng p-nitroaniline released / (min · mg protein)	338 ± 46 Q = 7.35§§§	540 ± 40 Q = 12.25***	456 ± 33 (42%) Q = 4.90***
Acid ribonuclease, change in absorbance / (min · mg protein)	0.043 ± 0.009 Q = 3.43§§§	0.065 ± 0.004 Q = 9.33***	0.052 ± 0.002 (62%) Q = 5.90***
Succinate dehydrogenase, change in absorbance / (min · mg protein)	0.0081 ± 0.0009 Q = 0.66 §	0.0123 ± 0.0016 Q = 6.94***	0.0086 ± 0.0007 (88%) Q = 6.12***
Glucose-6-phosphatase, nmol of P <sub>i</sub> released / (min · mg protein)	82 ± 6 Q = 0.46 §	47 ± 8 Q = 13.56***	81 ± 3 (97%) Q = 13.10***
Cytochrome P-450, pmol / (mg microsomal protein)	386 ± 68 Q = 9.93§§§	86 ± 61 Q = 11.92***	144 ± 56 Q = 1.98 <sup>+</sup>
Glycogen, mg · g <sup>-1</sup>	78 ± 11 Q = 25.18§§§	1.97 ± 0.25 Q = 26.55***	5.9 ± 4.3 Q = 1.37 <sup>+</sup>

(68%), γ-glutamyl transpeptidase (42%), acid ribonuclease (62%), and succinic dehydrogenase (88%). Almost complete restoration of the reduced activity of glucose-6-phosphatase was also noted. However, the decreased content of cytochrome P-450 and glycogen in liver remained unaffected by Pic.

#### DISCUSSION

Rats are quite refractory to α-amanitin and they develop phalloidin-like toxicity when poisoned with whole mushroom<sup>(9)</sup>. Phalloidin acts at the membrane by binding to actin-like protein, thus causing damage of membrane integrity<sup>(10)</sup>. Our results also showed that AP increased the activities of enzymes 5'-nucleotidase and γ-glutamyl transpeptidase as consequence of damage to plasma membrane of liver cells. Significant protection in both of these enzymes was also observed after Pic administration. We have earlier reported that

Pic provided significant restoration in the biochemical changes induced by D-galactosamine<sup>(2)</sup>. In this case the structural and functional impairment in membrane is perhaps one of the mechanisms involved. Histological studies<sup>(11)</sup> showed that phalloidin acts primarily on the membranes of liver lysosomes leading to release of lysosomal enzymes. Our results also showed increased activity of acid ribonuclease following administration of AP. Its reversal by Pic indicated that the damage of lysosomal membrane caused by AP is repaired by Pic administration. The alterations in the activities of succinic dehydrogenase, glucose-6-phosphatase and cytochrome P-450 by AP is indicative of altered mitochondrial, microsomal, and drug metabolizing functions of the liver. The activities of succinic dehydrogenase and glucose-6-phosphatase have significantly been restored by Pic, thus demonstrated its effectiveness against the

mitochondrial and microsomal damages. The drastic decrease in the levels of cytochrome P-450 and glycogen in liver remained unaffected with Pic, and a longer exposure of Pic might be required. Experiments are proceeding to elucidate the exact mechanism of action of Pic in antagonising the effects of AP and other toxic agents on liver.

**ACKNOWLEDGMENTS** We are grateful to Dr D K Kulshrestha for preparing Pic. AP lyophilysate was kindly provided by Prof G L Floersheim, Switzerland. Technical assistance by N K Verma and Suresh Yadav is also acknowledged.

## REFERENCES

- 1 Dwivedi Y, Rastogi R, Chander R, Sharma SK, Kapoor NK, Garg NK, *et al.* Hepatoprotective activity of picroliv against carbon tetrachloride-induced liver damage in rats. *Indian J Med Res* [B] 1990; **92** : 195-200.
- 2 Dwivedi Y, Rastogi R, Garg NK, Dhawan BN. Picroliv and its components Kutkoside and Picoside I protect liver against galactosamine-induced damage in rats. *Pharmacol Toxicol.* In press.
- 3 Dwivedi Y, Rastogi R, Sharma SK, Mehrotra R, Garg NK, Dhawan BN. Picroliv protects against monocrotaline-induced hepatic damage in rats. *Pharmacol Res* 1991; **23** : 399-407.
- 4 Dwivedi Y, Rastogi R, Garg NK, Dhawan BN. Prevention of paracetamol-induced hepatic damage in rats by picroliv, the standardized active fraction from *Picrorhiza kurroa*. *Phytotherapy Res* 1991; **5** : 115-9.
- 5 Dwivedi Y, Rastogi R, Sharma SK, Garg NK, Dhawan BN. Picroliv affords protection against thioacetamide-induced hepatic damage in rats. *Planta Med* 1991; **57** : 25-8.
- 6 Chander R, Dwivedi Y, Rastogi R, Sharma SK, Garg NK, Kapoor NK, *et al.* Evaluation of hepatoprotective activity of picroliv (from *Picrorhiza kurroa*) in *Mastomys natalensis* infected with *Plasmodium berghei*. *Indian J Med Res* [B] 1990; **92** : 34-7.
- 7 Govindan VM, Faulstich H, Wieland T, Agostini B, Hasselbach W. *In vitro* effect of phalloidin on a plasma membrane preparation from rat liver. *Naturwissenschaften* 1972; **59** : 521-2.
- 8 Floersheim GL, Bieri A, Koenig R, Pletsch A. Protection against *Amanita phalloides* by the iridoid glycoside mixture of *Picrorhiza kurroa* (Kutkin). *Agents Actions* 1990; **29** : 386-7.
- 9 Baccino FM, Cantino D, Zuretti MF. Studies on the hepatotoxicity of *Amanita phalloides* in the rat. I. Liver cell vacuolation. *Exp Mol Pathol* 1976; **24** : 159-75.
- 10 Frimmer M, Kroker R. The role of extracellular  $K^+$ -concentration in phalloidin effects on isolated hepatocytes and perfused rat livers. *Naunyn-Schmiedebergs Arch Pharmacol Exp Pathol* 1973; **278** : 285-92.
- 11 Rita GA, Zuretti MF, Baccino FM, Dianzani MU. The behaviour of liver lysosomes in *Amanita phalloides*. In: Giordano A, Zanbotti V, Matturri L, editors. *Enzyme histochemistry Symposium*. A. Baselli, Milan 1967; 1: 3-20.

## Instructions to authors

Please read *Acta Pharmacol Sin* 1992 Jan; **13**(1) : 3-8.

*Br Med J* 1991 Feb 9; **302**(6772) : 388-41.

*N Engl J Med* 1991 Feb 7; **324**(6) : 424-8.