

- 6 Mikkelsen EO, Thastrup O, Christensen SB. Effects of thapsigargin in isolated rat thoracic aorta. *Pharmacol Toxicol* 1988; 62 : 7-11.
- 7 Thastrup O. Role of Ca^{2+} -ATPase in regulation of cellular Ca^{2+} signalling, as studied with the selective microsomal Ca^{2+} -ATPase inhibitor, thapsigargin. *Agents Actions* 1990; 29 : 8-15.
- 8 Rees DD, Palmer RMJ, Schulz R, Hodson HF, Moncada S. Characterization of three inhibitors of endothelial nitric oxide synthase *in vitro* and *in vivo*. *Br J Pharmacol* 1990; 101 : 746-52.
- 9 Kwan CY, Deng HW, Guan YY. Tetrandrine is not a selective calcium channel blocker in vascular smooth muscle. *Acta Pharmacol Sin* 1992; 13 : 385-90.
- 10 Foskett JK, Wong D. Calcium oscillations in parotid acinar cells induced by microsomal Ca^{2+} -ATPase inhibition. *Am J Physiol* 1992; 262 : C656-63.
- 11 Bourreau JP, Abela AP, Kwan CY, Daniel EE. Acetylcholine Ca^{2+} stores refilling directly involves a dihydropyridine-sensitive channel in dog trachea. *Am J Physiol* 1991; 261 : C497-505.
- 12 Palmer RMJ, Ashton DS, Moncada S. Vascular endothelial cells synthesize nitric oxide from L-arginine. *Nature* 1988; 333 : 664-6.

BIBLID: ISSN 0253-9756 中国药理学报 *Acta Pharmacologica Sinica* 1993 Jan; 14 (1) : 26-30

Platelet activating factor production in bovine cerebral microvascular endothelial cells and its drug inhibition

SUN Du-Xin, HUANG Tong-Ge, ZENG Guo-Qian, RUI Yao-Cheng
(Department of Pharmacology, School of Pharmacy, Second Military Medical University, Shanghai 200433, China)

ABSTRACT The production of platelet activating factor (PAF) in bovine cerebral microvascular endothelial cells (CME cells) and the effects of tetrandrine (Tet) and dauricine (Dau) on the PAF production were investigated. PAF was determined by the aggregation of washed rabbit platelets. The results showed that the CME cells produced PAF $5.93 \text{ ng} / 8.5 \times 10^5 \text{ cells}$ under the calcimycin $2.5 \mu\text{mol} \cdot \text{L}^{-1}$ stimulation. Tet and Dau 1, 10, and 100 $\mu\text{mol} \cdot \text{L}^{-1}$ inhibited the production of PAF by 18.2%, 51.8%, 56.8%, and 26.3%, 63.3%, 65.9%, respectively. Tet concentration-dependently inhibited the PAF $9.1 \text{ nmol} \cdot \text{L}^{-1}$ induced washed rabbit platelets aggregation with the IC_{50} of $3.05 \mu\text{mol} \cdot \text{L}^{-1}$ (95% confidence limits: $0.59-15.86 \mu\text{mol} \cdot \text{L}^{-1}$). The binding of [^3H]triazolodiazepine to the CME cells was partially displaced by Tet $0.02-33.00 \mu\text{mol} \cdot \text{L}^{-1}$. It is suggested that the cerebrovascular system produces PAF at the pathological conditions and the inhibition of Tet and Dau.

KEY WORDS tetrandrine; dauricine; platelet activating factor; vascular endothelium; capillaries; triazoles.

Platelet activating factor (PAF) was originally described as a potent platelet activator derived from antigen-stimulated, IgE-sensitized rabbit basophils⁽¹⁾. PAF is involved in various disorders, such as asthma, shock, cardiovascular disorders, and renal ischemia⁽²⁾. PAF is 1000 to 10000 times more potent on a molar basis than histamine in increasing the vascular permeability. It is produced by various cells including basophils, macrophages, monocytes, and mast cells. PAF can be released from endothelial cells of human umbilical veins, bovine aortas, and pulmonary arteries^(3,4). It constricts the pial arteries of newborn pigs⁽⁵⁾, and decreases the cerebral blood flow of rats⁽⁶⁾. Our previous studies showed that the PAF specific binding

Received 1991-10-03

Accepted 1992-09-09

sites were existence on CME cells⁽⁷⁾. But the production of PAF in the cerebral microvascular endothelial cells (CME cells) has not been described. In the present study, we investigated the production of PAF and the inhibitory effects of Tet and Dau in CME cells.

MATERIALS AND METHODS

Calcimycin (A-23187, Sigma); trypsin (Sigma); [³H]triazolodiazepine (Tri) and C₁₆-PAF (Boehringer-er); Minimum essential medium (MEM, Gibco); BN 50739 (Gifted from P Braquet); Tris-Tyrodé's bovine serum albumin (BSA): pH 7.4, KCl 2.62, MgCl₂·6H₂O 1.0, NaCl 137, CaCl₂·2H₂O 1.30, Tris 10, glucose 5.6 mmol·L⁻¹, BSA 0.25%. Tyrodé's gelatin no calcium (TG-no calcium): pH 6.5, KCl 2.62, MgCl₂·6H₂O 1.0, NaCl 137, NaHCO₃ 12, glucose 5.6, EGTA 0.2 mmol·L⁻¹, gelatin 0.25%. Tris-HCl buffer: pH 7.4, KCl 140 mmol·L⁻¹, Tris 10 mmol·L⁻¹, BSA 0.1%. Acid citrate dextrose: citric acid 6.0, sodium citrate 8.5, glucose 12.3 mmol·L⁻¹.

Cultivation of bovine CME cells and PAF production Cells were cultivated by our previous method⁽⁸⁾. The CME cells were collected with 0.1% trypsin from culture flask, washed with Tris-Tyrodé's BSA (800× g for 8 min), resuspended in 0.5 ml Tris-Tyrodé's BSA (1.7×10⁶ cells·ml⁻¹), and incubated 10 min at 37°C. After Tet and Dau (1-100 μmol·L⁻¹) were added, the cells were incubated for another 10 min further. Finally, the cells were stimulated with calcimycin (2.5 μmol·L⁻¹) for 20 min. Then the suspension was acidified with 20 μl acetic acid. PAF was extracted⁽⁹⁾. The CHCl₃ phase were collected by centrifugation of 1200× g for 15 min and dried by the stream of N₂. The residues were kept at -40°C before use.

Aggregation of rabbit platelet rich plasma (PRP) and washed platelets Blood was collected from the heart of New Zealand white rabbits (3.2±0.3 kg) with plastic syringes containing acid citrate dextrose (1 to 6 vol of blood). Platelet-rich plasma (PRP) was prepared by centrifugation of 400× g for 6 min.

Platelets were obtained by centrifugation of 1400× g for 15 min. After gently washed with TG-no calcium (1400× g for 15 min), the platelets were resuspended in TG-no calcium containing acetylsalicylic acid (ASA) 0.1 mmol·L⁻¹ for 15 min at 25°C, and centrifuged (1400× g for 15 min). The platelets were gently suspended in Tris-Tyrodé's BSA at a concentration of 4×10⁸ platelets·ml⁻¹. The aggregation of platelets were induced by PAF. Creatine phosphate-creatine phosphokinase enzymatic system (CP/CPK 312.5/152.5 μg·ml⁻¹) was directly added to the platelets 30 s before PAF was added. The platelets aggregation of PRP were induced by adenosine diphosphate (ADP) 5 μmol·L⁻¹, or arachidonic acid (AA) 80 μmol·L⁻¹, or PAF 9.1 nmol·L⁻¹.

Assay of PAF PAF was assayed by the aggregation of washed rabbit platelets in an aggregometer. Extracts of the CME cells were dissolved in Tris-Tyrodé's BSA, and 10 μl were added to the platelets. Platelet aggregation induced by extracts was recorded, and the amount of PAF was calculated on a calibration curve of platelet aggregation induced by synthetic PAF (0.27-9.1 nmol·L⁻¹).

PAF binding assay The cells were collected with 0.1% trypsin, resuspended in Tris-HCl buffer (6×10⁵ cells·ml⁻¹). The cell suspension (3×10⁵ cells) was added to a tube containing [³H]Tri 8 nmol·L⁻¹. Tet, with or without 1000 fold unlabeled Tri, was incubated for 15 min at 25°C. Bound and free radioligands were separated by rapid filtration with microporous filtering film (0.8 μm). The filters were washed 5 times with ice-cold Tris-HCl buffer (no BSA). The radioactivity was measured by scintillation.

RESULTS

PAF activity in CME cells In washed rabbit platelets, the action of arachidonic acid and ADP were blocked by acetylsalicylic acid and CP/CPK. ADP 5 μmol·L⁻¹ or arachidonic acid 80 μmol·L⁻¹ induced the aggregation of PRP, but not the aggregation

of the washed platelets at the same concentration (Tab 1). The aggregations of PRP and the washed platelets were induced by PAF 0.27–9.1 nmol·L⁻¹, and blocked by PAF receptor antagonist Tri 10 μmol·L⁻¹. But Tri did not block the platelet aggregation of PRP induced by ADP or arachidonic acid (Tab 1). The aggregation of washed platelets was induced by the extracts of CME cells and blocked by Tri 10 μmol·L⁻¹ (Tab 1).

Tab 1. Aggregation (%) of platelet-rich plasma (PRP) and washed rabbit platelets (WRP) induced by arachidonic acid (AA), ADP, platelet activating factor (PAF), and the antagonism of triazolodiazepine (Tri) 10 μmol·L⁻¹. n=3 rabbits. $\bar{x} \pm s$.

Stimulants	PRP	WRP
AA 80 μmol·L ⁻¹	83±8	0
ADP 5 μmol·L ⁻¹	60±7	0
PAF 9.1 nmol·L ⁻¹	66±7	65±6
EXtracts of CME cells (4×10 ⁴)	45±8	44±10
AA 80 μmol·L ⁻¹ + Tri	80±8	0
ADP 5 μmol·L ⁻¹ + Tri	58±7	0
PAF 9.1 nmol·L ⁻¹ + Tri	0	0
Extracts of CME cells (4×10 ⁴ + Tri)	0	0

PAF production induced by calcimycin

PAF was produced by CME cells under the stimulation of calcimycin, but it was not synthesized without calcimycin stimulation. The amount of PAF attained the peak at calcimycin 2.5 μmol·L⁻¹ after 20 min incubation, and it decreased gradually after further stimulation (Tab 2).

Inhibitory effects of Tet and Dau on production of PAF in CME cells CME cells produced PAF 5.93 ng/8.5×10⁵ cells under the stimulation of calcimycin 2.5 μmol·L⁻¹ for 20 min. Tet and Dau 1, 10, and 100

Tab 2. Stimulation of PAF production by calcimycin 2.5 μmol·L⁻¹ for 20 min. n=3, $\bar{x} \pm s$. **P<0.05 vs 10 min group, *P<0.01 vs 20 min group, a: P<0.01 vs calcimycin 0.1 μmol·L⁻¹ group, b: P<0.01 vs calcimycin 1.0 μmol·L⁻¹ group, c: P<0.01 vs calcimycin 2.5 μmol·L⁻¹ group with Newman-Keuls test.**

Time, min	PAF, ng/8.5×10 ⁵ cells	Calcimycin, μmol·L ⁻¹	PAF, ng/8.5×10 ⁵ cells
10	0.97±0.21	0.1	0.39±0.10
20	5.08±0.78**	1.0	1.02±0.29
45	2.21±0.38***	2.5	6.04±1.20 ^{a,b}
60	1.60±0.52***	10.0	2.85±0.36 ^{a,b,c}

μmol·L⁻¹ incubated for 10 min with CME cells inhibited the production of PAF by 18.2%, 51.8%, 56.8%, and 26.3%, 63.3%, 65.9%, respectively (Fig 1).




Fig 1. Inhibition of tetrandrine (Tet) and dauricine (Dau) on PAF production in cerebral microvascular endothelial cells (CME cells). n=5, $\bar{x} \pm s$. *P>0.05, **P<0.05, *P<0.01 vs control with Dunnett test.**

Inhibition of Tet on aggregation of washed platelets induced by PAF Tet 0.4135, 4.135, 41.35, and 82.7 μmol·L⁻¹ or PAF receptor antagonist BN 50739 1, 10, 100, and 1000 nmol·L⁻¹ preincubated for 1 min with platelets concentration-dependently inhibited PAF induced platelet aggregation by 24.6%,

45.9%, 86.9%, 100%, and 1.6%, 32.3%, 88.7%, 100%, respectively, their IC_{50} were $3.05 \mu\text{mol} \cdot \text{L}^{-1}$ (95% confidence limits: $0.59-15.86 \mu\text{mol} \cdot \text{L}^{-1}$) and $18.73 \text{nmol} \cdot \text{L}^{-1}$ (95% confidence limits: $4.23-82.17 \text{nmol} \cdot \text{L}^{-1}$), respectively (Fig 2).




Fig 2. Effects of Tet and BN 50739 on aggregation of washed rabbit platelets induced by PAF. $n=3$, $\bar{x} \pm s$. * $P > 0.05$, ** $P < 0.05$, *** $P < 0.01$ vs control with Dunnett test.

Inhibition of Tet on the $[^3\text{H}]\text{Tri}$ binding to CME cells The specificity of $[^3\text{H}]\text{Tri}$ binding to CME cells was established by displacing $[^3\text{H}]\text{Tri}$ $8 \text{nmol} \cdot \text{L}^{-1}$ by Tet in competition experiment. The binding of $[^3\text{H}]\text{Tri}$ was partially displaced by Tet ($0.017-33.00 \mu\text{mol} \cdot \text{L}^{-1}$) (Tab 3).

DISCUSSION

PAF could be produced by bovine and rat brain. The existence of PAF in the brain was characterized by GC-MS, HPLC, and TLC, quantified the amount of PAF by bioassay^(10,11). Our results firstly showed that the CME cells synthesized PAF $5.93 \text{ng} / (8.5 \times 10^5 \text{ cells})$ under calcimycin $2.5 \mu\text{mol} \cdot \text{L}^{-1}$ stimulation. But human umbilical cord veins endothelial cells produced PAF $21.5 \text{ng} / (5 \times 10^5 \text{ cells})$ at calcimycin $2.5 \mu\text{mol} \cdot \text{L}^{-1}$ stimulation⁽¹²⁾. This might be the difference between the large blood vessels and the cere-

bral microvessels. The amount of PAF reached maximum after 20 min of stimulation and decreased thereafter gradually. This might be due to a rapid increase in the acetyltransferase activity and an unchange-ment of acetylhydrolase activity⁽¹³⁾

Tab 3. Inhibition of tetrandrine (Tet) on $[^3\text{H}]\text{Tri}$ binding to cerebral microvascular endothelial cells (CME cells). $n=3$, $\bar{x} \pm s$. * $P > 0.05$, ** $P < 0.05$ vs control with Dunnett test.

Tet, $\mu\text{mol} \cdot \text{L}^{-1}$	Bound, $\text{dpm} / 3 \times 10^5 \text{ cells}$	Inhibition, %
Control	$22\ 318 \pm 2\ 032$	
0.02	$18\ 716 \pm 1\ 892^*$	16.1
0.17	$14\ 494 \pm 2\ 838^*$	35.1
1.65	$11\ 870 \pm 1\ 378^{**}$	46.8
33.00	$10\ 562 \pm 943^{**}$	52.7

In the bioassay of PAF, the extracts of CME cells was added directly to the washed rabbit platelets. The aggregation of platelets induced by the extracts could be blocked by PAF receptor antagonist Tri. It is similar to the reaction of synthetic PAF. These suggested that the activity of the extracts was that of PAF.

Tet and Dau inhibited the production of PAF. Tet inhibited the specific binding of $[^3\text{H}]\text{Tri}$ to CME cells and the aggregation of washed rabbit platelets induced by PAF. Our previous studies showed that Tet and Dau inhibited the proliferation and the DNA synthesis of bovine cerebral microvascular smooth muscle cells induced by PAF. Dau inhibited the binding of $[^3\text{H}]\text{Tri}$ to the cerebral microvascular smooth muscle cells. It suggested that Tet and Dau might inhibit the PAF action in cerebral vascular system.

REFERENCES

1 Benveniste J, Henson PM, Cochrane CG. Leukocyte-dependent histamine release from rabbit platelets. The role of IgE, basophils, and a platelet-activating factor. *J Exp Med* 1972; 136 : 1356-77.

2 Benveniste J. PAF-acether, an ether phospho-lipid with biological activity. *Prog Clin Biol Res* 1988; 282 : 73-85.

3 McIntyre TM, Zimmerman GA, Satoh K, Prescott SM. Cultured endothelial cells synthesize both platelet-activating factor and prostacyclin in response to histamine, bradykinin, and adenosine triphosphate. *J Clin Invest* 1985; 76 : 271-80.

4 Whatley RE, Zimmerman GA, McIntyre TM, Prescott SM. Endothelium from diverse vascular sources synthesizes platelet-activating factor. *Arteriosclerosis* 1988; 8 : 321-31.

5 Armstead WM, Pourcyrous M, Mirro R, Leffler CW, Busija DW. Platelet activating factor: A potent constrictor of cerebral arterioles in newborn pigs. *Circ Res* 1988; 62 : 1-7.

6 Kochanek PM, Nemoto EM, Melick JA, Evans RW, Burke DF. Cerebrovascular and cerebro-metabolic effects of intracarotid infused platelet-activating factor in rats. *J Cerebral Blood Flow Metab* 1988; 8 : 546-51.

7 Sun DX, Rui YC, Zeng GQ, Zhu J, Shen YA. Specific Binding sites of platelet activating factor on the intact bovine cerebral microvascular endothelial cells and antagonism of drugs. *Acta Pharmacol Sin* 1992; 13 : 326-9.

8 Sun DX, Zeng GQ, Shen YA, Rui YC. Cultivation of bovine cerebral microvascular endothelial cells in vitro. *Acad J Sec Mil Med Univ* 1992; 13 : 178-80.

9 Bligh EG, Dyer WJ. A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* 1959; 37 : 911-7.

10 Tokumura A, Kamiyasu K, Takauchi K, Tsukatani H. Evidence for existence of various hemologues and analogues of platelet activating factor in a lipid extract of

bovine brain. *Biochem Biophys Res Commun* 1987; 145 : 415-25.

11 Kumar R, Harvey SAK, Kester M, Hanahan DJ, Olson MS. Production and effects of platelet-activating factor in the rat brain. *Biochim Biophys Acta* 1988; 963 : 375-83.

12 Camussi G, Aglietta M, Malavasi F, Tetta C, Piacibello W, Sanavio F, et al. The release of platelet-activating factor from human endothelial cells in culture. *J Immunol* 1983; 131 : 2397-403.

13 Hirafuji M, Mencia-Huerta JM, Benveniste J. Regulation of PAF-acether (platelet-activating factor) biosynthesis in cultured human vascular endothelial cells stimulated with thrombin. *Biochim Biophys Acta* 1987; 930 : 359-69.

26-30

(3)

血小板激活因子在脑微血管内皮细胞上的产生及药物的抑制作用

R 973

孙笃新、黄彤莉、曾国钱、芮耀诚 (第二军医大学药学院药理教研室, 上海 200433, 中国)

摘要 在体外培养的脑微血管内皮细胞上研究血小板激活因子(PAF)的产生。结果表明, 该细胞在卡西霉素刺激下可产生大量 PAF, 粉防己碱和蝙蝠葛碱对 PAF 的产生具有抑制作用, 而且粉防己碱能够抑制 PAF 诱导的兔洗涤血小板的聚集和 [³H]triazolodiazepine 与内皮细胞的结合。结果提示, 粉防己碱能够抑制 PAF 的作用。

PAF

关键词 粉防己碱; 蝙蝠葛碱; 血小板激活因子; 血管内皮; 微血管; 三唑类

国家医药和农药筛选协作组征集样品启事

由中国科学院上海药物研究所, 北京医科大学, 中国医学科学院药物研究所和南开大学四个单位组成的医药和农药筛选协作组已正式成立, 并向全国开放, 免费筛选各种医用和农用化合物, 评价其生物活性。欢迎全国各地向协作组寄送待测样品。联系地址: (医用化合物) 200031 上海市太原路 294 号中科院上海药物所筛药办公室; 100083 北京市学院路北京医科大学天然药物及仿生药物国家重点实验室; 100050 北京市先农坛街 1 号中国医科院药物研究所计划处。(农用化合物) 300071 天津市卫津路 94 号天津南开大学元素有机化学国家重点实验室。