

Prostacyclin-thromboxane interactions in the platelet-perfused in vitro heart.
Am J Physiol 1981; **241** : H18-25.

4 Alloatti G, Montrucchio G, Camussi G. Prostacyclin inhibits platelet-dependent effects of the platelet activating factor in the isolated rabbit heart.
J Cardiovasc Pharmacol 1990; **15** : 745-51.

5 Ardlie NG, Packham MA, Mustard JF. Adenosine diphosphate-induced platelet aggregation in suspensions of washed rabbit platelets.
Br J Haematol 1970; **19** : 7-17.

6 Tang XL, Guo ZG. The effect of EDTA on perfused isolated working guinea pig hearts.
Bull Hunan Med Col 1987; **12** : 125-8.

7 Rönberg AL, Hansson C, Håkanson R. High-performance liquid chromatographic determination of histamine in biological samples after derivation with *o*-phthalaldehyde.
Anal Biochem 1984; **139** : 338-44.

8 Xie DN, Chen YL, Guo ZG. A fast and reliable OPA derivatization method to determine histamine concentration in washed rabbit platelet suspension by HPLC.
Chin J Chromatogr. In press.

9 Ashmore SP, Thomson AH, Simpson H. High-performance liquid chromatographic technique for the rapid determination of histamine in both the plasma and cellular components of blood.
J Chromatogr 1989; **496** : 435-40.

10 Levi R, Owen DAA, Trzeciakowski J. Actions of

histamine on the heart and vasculature. In: Ganellin CR, Parsons ME, editors. *Pharmacology of histamine receptors*. Bristol: John Wright, 1982; 236-67.

11 Martin JF, Trombridge EA, Salmon GL, Slater DN. The relationship between platelet and megakaryocyte volumes. *Thromb Res* 1982; **28** : 447-59.

12 Saxena SP, Brandes LJ, Becker AB, Simons KJ, LaBella FS, Gerrard JM. Histamine is an intracellular messenger mediating platelet aggregation.
Science 1989; **243** : 1596-9.

93,14(5)
385-388

家兔血小板对离体豚鼠作功心脏的增强效应及组胺的参与

郭兆贵, 陈亚立, 谢大年 R965.2
(湖南医科大学药理研究室, 长沙410078, 中国)

摘要 用洗涤家兔血小板悬液和其上清灌注离体豚鼠作功心脏, 心脏作功浓度依赖性增强。西咪替丁(Cim) 1 $\mu\text{mol}\cdot\text{L}^{-1}$ 显著抑制增强相。经 HPLC 测出血小板悬液每 10^9 血小板含组胺 $2.6 \pm 0.7 \mu\text{g}$ 。电镜显示受血小板攻击的心脏肥大细胞处于活动期。结果表明, 心功能加强与血小板及心脏肥大细胞释放的组胺有关。

关键词 组胺; 西咪替丁; 血小板; 心脏; 血液动力学; 肥大细胞; 电镜

Chlorpromazine attenuated electroacupuncture analgesia in conscious rabbits¹

DAI Jia-Le, XU Shao-Fen (Department of Neurobiology, School of Basic Medical Sciences, Shanghai Medical University, Shanghai 200032, China)

ABSTRACT By measuring the defense behavior in response to the noxious stimulation induced by potassium iontophoresis on ear-lobe skin of conscious rabbit, chlorpromazine (CPZ) ($0.5 \text{ mg}\cdot\text{kg}^{-1}$, iv) induced hyperalgesia, whereas it significantly attenuated electroacupuncture analgesia (EAA) efficacy. Mono-

amines and their metabolites in cerebrospinal fluid (CSF) were measured by high pressure liquid chromatography with electrochemical detector (HPLC-ECD) while the attenuation effect of CPZ on EAA was observed. CPZ markedly enhanced 3,4-dihydroxyphenylacetic acid (DOPAC) ($P < 0.05$) and homovanillic acid (HVA) ($P < 0.01$) contents in CSF both in the presence and absence of electroacupuncture. CPZ attenuated EAA with elevations of either DOPAC or HVA concentration in CSF. There was a positive cor-

Received 1991-09-26 Accepted 1993-02-17

¹ Presented at the 4th National Symposium on Acupuncture Anesthesia & Acupuncture Analgesia, 1991 Sep 17-20, Xi-an, China.

relation between the increase of DOPAC or HVA content in CSF and the attenuation effect of CPZ on EAA ($P < 0.05$). These results suggested that the activation of dopamine system was unfavorable for EAA.

KEY WORDS electroacupuncture; analgesia; chlorpromazine; dopamine receptors; 3, 4-dihydroxyphenylacetic acid; homovanillic acid; cerebrospinal fluid; biogenic monoamines; high pressure liquid chromatography

Monoamines^[1-3], endogenous opioid peptides (EOP)^[4,5], and acetylcholine played important roles in electroacupuncture analgesia (EAA), which altered the levels of various neurotransmitters in the central nervous system (CNS). In previous experiments, involvement of dopaminergic mechanisms in EAA has been demonstrated^[2,3], and dopamine (DA) receptor antagonists were consequently categorized as EAA synergists^[3,6]. Chlorpromazine (CPZ), a neuroleptic mainly due to its anti-dopaminergic action, weakened EAA in rabbits^[7]. CPZ possesses a broad spectrum of pharmacological actions, especially in modulating the actions of monoamine transmitters in CNS, i.e., anti-DA, anti-serotonin (5-HT), anti-norepinephrine (NE), etc^[6]. To probe into the intrinsic mechanism underlying the attenuation effect of CPZ on EAA, we observed the effect of CPZ on EAA, and simultaneously determined the content changes of monoamines and their metabolites in cerebrospinal fluid (CSF) of rabbits after CPZ and/or electroacupuncture (EA).

MATERIALS AND METHODS

CPZ was purchased from Shanghai Tianfeng Pharmaceutical Factory. NE, 3-methoxy-4-hydroxyphenylglycol (MHPG), DA, 3, 4-dihydroxyphenylacetic acid (DOPAC), homovanillic acid (HVA), 5-HT, and 5-hydroxyindoleacetic (5-HIAA) were the products from Sigma Co.

New Zealand rabbits (bred by Shanghai Medical

University) of either sex, weighing 2.1 ± 0.2 kg, were used.

Rabbit skin pain model The potassium iontophoresis method was used to measure the pain threshold in conscious rabbits. K^+ were brought into the ear-lobe skin by continuously increasing direct current as noxious stimulus, and withdrawal reactions of forelimbs and/or head were served as a criterion of pain response, while the current value (mA) was recorded as pain threshold. Before administration of drug and/or EA, the pain threshold of each rabbit was measured 3 times at 5-min intervals, and the \bar{x} was used as the baseline. Rabbits with pain thresholds > 1 mA were discarded. Pain thresholds were tested at 10-20-min intervals during 100 min following administration.

Electroacupuncture EA was carried out at "Hegu" point (midpoint of medial edge of 2nd metacarpal of forepaw) and "Waiguan" point (dorsum of forelimb, 12 mm above wrist joint along the medium line between radius and ulna). Unilateral points were electrically needled for 30 min by EA apparatus (Model G-6805-2), at a frequency of 3 Hz with an intensity capable of inducing slight tremble of forelimb. Sham needling (SN) without electric stimulation was used as control.

Cannula implantation and CSF collection Rabbits were anesthetized with sodium pentobarbital ($30 \text{ mg} \cdot \text{kg}^{-1}$, iv). A plastic cannula (OD=0.9 mm, ID=0.6 mm) was implanted into the 4th ventricle (coordinates P 13, L/R 0.5, H 0, according to Sawyer's Atlas). The cannula was then clogged up and secured by dental cement. Rabbits were individually caged until they were tested 3 d later. At the beginning and 30th min following administration, pain thresholds were recorded while CSF ($30 \mu\text{l}$) was collected within 30 s. The CSF were collected in tubes containing $10 \mu\text{l}$ HCl ($0.1 \text{ mol} \cdot \text{L}^{-1}$), frozen with dry ice, and stored at -30°C until assay.

High pressure liquid chromatography coupled to electrochemical detector (HPLC-ECD) Monoamines and their metabolites were detected by ion-pair reverse phase HPLC-ECD. The HPLC-ECD system (Waters) consisted of a M510 pump, a U6K injection system coupled to a M460 electrochemical detector and M740 data module. The separation was accomplished on a μ -Bondapak C_{18} column (300×3.9 mm ID) with a pre-

column (Guard Pak TMRCS C₁₈). The eluent consisted of chloro-acetic acid-sodium hydroxide buffer 0.15 mol·L⁻¹ containing EDTA 0.83 mmol·L⁻¹, D-camphor-β-sulfonic acid 9 mmol·L⁻¹, and 5% methanol (pH=4.2), at a flow rate of 1.5 ml·min⁻¹. The eluent was filtered through a 0.22 μm membrane and degassed before use. The working potential was +0.7 V. Samples of CSF were injected directly.

Protocol Rabbits were randomly divided into 4 groups: (1) NS+SN group; (2) CPZ+SN group; (3) EA+NS group; (4) EA+CPZ group. Results were expressed as $\bar{x} \pm s$, and statistical significance between groups was evaluated by *t* test.

RESULTS

Effect of CPZ on pain threshold and EAA
 Effect of CPZ on baseline pain threshold as well as EAA was examined on 32 rabbits. The rabbits elicited hypersensitivity to pain after CPZ (0.5 mg·kg⁻¹, iv) treatment. Compared with NS+SN control group, the pain threshold was significantly lowered. Given concurrently with EA, such a dose of CPZ weakened the EAA efficiency (Tab 1). The pain threshold change of EA+CPZ group was less than that of the sum of CPZ group and EA group, ie, CPZ markedly antagonized EAA. The attenuation action maintained for 80 min, the pain threshold remained little enhanced or even lower than the preadministration level during 100 min following administration.

Influence of CPZ and/or EA on monoamine levels in CSF NE, DA, MHPG, DOPAC, HVA, and 5-HIAA, but not 5-HT were detectable in CSF of rabbits, and the baseline monoamine contents of metabolites (50–300 ng·ml⁻¹) were higher than their prototypes (0–10 ng·ml⁻¹), owing to the rapid metabolic rate in the CNS. The NE and MHPG contents remained unchanged after CPZ and/or EA treatment. There was no significant change of DA content in all the 4 groups. But levels of DOPAC and HVA, DA metabolites, were significantly increased after CPZ injection. EA induced no changes of DOPAC or HVA contents. When EA was applied in combination with CPZ, DOPAC, and HVA levels were augmented. EA brought forth a marked elevation of 5-HIAA level. When CPZ was administrated together with EA, the 5-HIAA content was decreased, but no statistical significance was shown (Tab 2).

Correlation between DOPAC or HVA levels and EAA effect CPZ attenuated the EAA effect with elevation of either DOPAC or HVA level, showing a positive correlation (Fig 1). This suggested that increase of DA metabolites in CSF was unfavorable for EAA.

Tab 1. Effect of CPZ (0.5 mg·kg⁻¹, iv) on pain threshold (mA) and EAA in conscious rabbits. *n* = 8, $\bar{x} \pm s$. Statistical significance between changes of pain thresholds was examined by *t* test. **P* > 0.05, ^b*P* < 0.05, ^c*P* < 0.01 vs NS+SN; ^d*P* > 0.05, ^e*P* < 0.05, ^f*P* < 0.01 vs NS+EA.

Time/min	NS+SN	CPZ+SN	NS+EA	CPZ+EA
0	0.70 ± 0.21	0.73 ± 0.12 ^a	0.67 ± 0.24 ^a	0.67 ± 0.17 ^d
10	0.72 ± 0.12	0.57 ± 0.23 ^b	1.25 ± 0.40 ^e	0.87 ± 0.40 ^e
20	0.69 ± 0.15	0.45 ± 0.20 ^c	1.61 ± 0.54 ^e	0.93 ± 0.34 ^f
30	0.70 ± 0.16	0.42 ± 0.31 ^c	1.36 ± 0.49 ^e	0.87 ± 0.48 ^e
40	0.67 ± 0.20	0.33 ± 0.35 ^c	1.17 ± 0.39 ^e	0.67 ± 0.50 ^e
60	0.68 ± 0.16	0.35 ± 0.31 ^c	0.98 ± 0.30 ^e	0.53 ± 0.45 ^f
80	0.73 ± 0.25	0.39 ± 0.31 ^b	0.82 ± 0.20 ^e	0.61 ± 0.26 ^e
100	0.69 ± 0.20	0.35 ± 0.30 ^c	0.73 ± 0.32 ^e	0.53 ± 0.25 ^d

Tab 2. Content changes of monoamines and their metabolites in CSF of conscious rabbits after CPZ and/or EA administration. $\bar{x} \pm s$. * $P > 0.05$, ^b $P < 0.05$, ^c $P < 0.01$ or NS+SN; ^d $P > 0.05$, ^e $P < 0.05$, ^f $P < 0.01$ or NS+EA.

(n)	Content increases of monoamines and their metabolites/ng·ml ⁻¹			
	NS+SN (8)	CPZ+SN (7)	NS+EA (8)	CPZ+EA (7)
NE	2.4±6.7	1.4±6.8 ^a	1.8±8.0 ^a	2.7±8.6 ^d
MHPG	1.3±8.8	0.8±9.8 ^a	5.8±11 ^a	6.5±11 ^d
DA	0.4±2.3	1.9±5.9 ^a	-0.6±2.3 ^a	-0.4±4.0 ^d
DOPAC	31±37	140±110 ^b	30±28 ^a	100±76 ^e
HVA	-30±78	180±180 ^c	-30±170 ^a	220±140 ^f
5-HIAA	-5.0±14	4.5±12 ^a	28±28 ^b	6.0±18 ^d




Fig 1. Correlation analysis between content increases of DOPAC and HVA in CSF of rabbits and attenuation rate of CPZ on EAA.

DISCUSSION

The hypothesis of the existence of two categories of DA receptors, D₁ and D₂, has been generally accepted⁽⁹⁾. It has been established that D₂, but not D₁ receptor, modulated the DA release and metabolism⁽¹⁰⁾, whereas DA presynaptic autoreceptors were in good accordance with D₂ type⁽⁹⁾. The antagonists for D₂ receptor potentially stimulated DA release, especially in lower dosage⁽¹¹⁾. CPZ, a potent DA receptor antagonist for both D₁ and D₂, had a preferential affinity for D₂ subtype. The present results showed a marked content elevation of DOPAC and HVA after CPZ (0.5 mg·kg⁻¹, iv) both in the presence and absence of EA. CPZ, used in low dosage in our study, blocked the presynaptic DA autorecep-

tors, activated the DA system, and thus provoked a content elevation of DA metabolites.

Studies concerning the influence of DA system on EAA has been intensively conducted. It was found that central dopaminergic activation was an unfavorable factor on EAA. Previous experiments indicated that DA receptor blockers consisted of a large family of EAA synergists, both from animal observations^(3,6) and in clinical therapy⁽⁷⁾. In addition, it has been revealed that the anti-dopaminergic action on CNS contributed to the potential effects of these drugs on EAA^(3,12). Our data analysis also revealed a positive correlation between the elevation of DOPAC or HVA content in CSF and the attenuation rate of CPZ on EAA. These studies suggested that CPZ attenuated EAA, probably via blocking the central DA autoreceptors, and thus activating the DA system. Furthermore, EAA was a complicated process involving the integration of EOP, biogenic monoamines, and other transmitter systems in CNS. Opiates modulated the synthesis, release and metabolism of biogenic amines⁽¹³⁾, and one-third of opioid receptors in striatum were present on the dopaminergic nerve terminals⁽¹⁴⁾, which strongly suggested the bilateral regulatory actions of DA and EOP system. Whether the attenuation effect of CPZ on EAA was EOP-mediated still remained unclear in this study.

There have been sound evidences showing that increases in the activity of brain and spinal cord 5-HT neurons are associated with analgesia⁽¹⁵⁾ as well as EAA⁽¹⁾. CPZ is considered to be a moderate 5-HT receptor blocker⁽⁹⁾. Consistant with previous experiments, our results revealed an increase of 5-HIAA content in CSF after EA application. We also found that 5-HIAA was lowered when CPZ was used in combination with EA, which confirmed the attenuation action of CPZ on EAA.

REFERENCES

- 1 Foster JM, Sweeney BP. The mechanisms of acupuncture analgesia. *Br J Hosp Med* 1987; 38 : 308-12.
- 2 Sun AY, Boney F, Lee DZ. Electroacupuncture alters catecholamines in brain regions of rats. *Neurochem Res* 1985; 10 : 251-8.
- 3 Xu SF, Cao XD, Mo WY, Xu ZB, Pan YY. Effect of combination of drugs with acupuncture on analgesic efficacy. *Acupunct Electrother Res* 1989; 14 : 103-13.
- 4 He LF. Involvement of endogenous opioid peptides in acupuncture analgesia. *Pain* 1987; 31 : 99-121.
- 5 Dai JL, Xu SF. Effect of tiapride on electroacupuncture analgesia. *Acta Pharmacol Sin* 1991; 12 : 212-5.
- 6 Xu ZB, Pan YY, Xu SF, Mo WY, Cao XD, He LF. Synergism between metoclopramide and electroacupuncture analgesia. *Acupunct Electrother Res* 1983; 8 : 283-8.
- 7 Dai JL, Cui YG, Zhang J, Xu SF, Wang HH, Jiang JW, et al. Attenuation of electro-acupuncture by chlorpromazine, chlorprothixene and promethazine. *N Drugs Clin Remed* 1990; 9 : 325-7.
- 8 Baldessarini RJ. Drugs and the treatment of psychiatric disorders. In: Gilman AG, Rall TW, Nies AS, Taylor P, editors. *The pharmacological basis of therapeutics*. NY: Pergamon, 1990 : 383-435.
- 9 Stoof JC, Keabian JW. Two dopamine receptors: bio-

chemistry, physiology and pharmacology.

Life Sci 1984; 35 : 2281-96.

- 10 Boyar WC, Altar CA. Modulation of *in vivo* dopamine release by D₂ but not D₁ receptor agonists and antagonists. *J Neurochem* 1987; 48 : 824-31.
- 11 Collu R, Bouvier C. Effects of sulpiride and apomorphine on prolactin release in adrenalectomized animals. Role of sodium ions. *Brain Res* 1987; 401 : 23-9.
- 12 Sun DY, Wang WQ, Jiang EK, Gao DM. Intrastantia nigra injection of dopamine on electroacupuncture analgesia in rabbits. *Acupunct Res* 1985; 10 : 30-2.
- 13 Gauchy C, Agid Y, Glowinski J, Cheramy A. Acute effects of morphine on dopamine synthesis and release and tyrosine metabolism in the rat striatum. *Eur J Pharmacol* 1973; 22 : 311-9.
- 14 Pollard H, Llorens-Cortes C, Schwartz JC. Enkephalin receptors on dopaminergic neurons in rat striatum. *Nature* 1977; 268 : 745-7.
- 15 Messing RB, Lytle LD. Serotonin-containing neurons: their possible role in pain and analgesia. *Pain* 1977; 4 : 1-21.

388-392

(2)

氯丙嗪在清醒家兔上使电针镇痛减效

戴佳乐, 许绍芬 (上海医科大学基础医学院神经生物学教研室, 上海200032, 中国)

R965.2

A

摘要 用钾离子透入法测定兔耳尖皮肤痛阈值, 观察氯丙嗪(CPZ)对痛阈及电针镇痛(EAA)的影响. CPZ (0.5 mg·kg⁻¹, iv)不仅降低痛阈值且减弱 EAA. 高压液相-电化学检测法测定发现 NE, MHPG, DA 含量在给予 CPZ 和或电针后均无明显改变. 而在 CPZ+伴针和+电针组中, DOPAC 和 HVA 含量均明显升高, 且升高值与 EAA 减效率呈正相关, 提示激活 DA 系统不利于 EAA.

关键词 电针; 镇痛; 氯丙嗪; 多巴胺受体; 3,4-二羟苯乙酸; 高香草酸; 脑脊液; 生物单胺类; 高压液相色谱法