

## REFERENCES

- 1 Wang Y, Li DX. Anti-arrhythmic action of *l*-tetrahydropalmatine. *Acta Pharmacol Sin* 1987; 8: 337-40.
- 2 Zhou J, Li DX. Cardiovascular pharmacology of *l*-tetrahydropalmatine and its analogues. *Chin Pharmacol Bull* 1992; 8: 253-6.
- 3 Zhao DH, Zhao GD, Sheng BH, Wang FC. The experimental anti-arrhythmic effect of *l*-stepholidine. *Chin Pharmacol Bull* 1988; 4: 35-8.
- 4 Ljima T, Irisawa H, Kameyama M. Membrane current and their modification by acetylcholine in isolated single atrial cells of the guinea pig. *J Physiol (Lond)* 1985; 359: 485-501.
- 5 Isenberg G, Klockner U. Glycocalyx is not required for slow inward calcium current in isolated rat heart myocytes. *Nature* 1980; 284: 258-60.
- 6 Chang TH, Li JM, Sun XD, Yu YF, Feng WY, Hao LY, et al. Effect of Pd-1a, a component of Chinese traditional herb Bai-Hua Qian-Hu, on  $Ca^{2+}$ -current and action potential duration in single ventricular cells of guinea pig hearts. *Shoyakugaku Zasshi* 1993; 47: 279-82.
- 7 Li JM, Chang TH, Sun XD, Hao LY, Wang YP, Yu YF, et al. Effect of dl-praeruptorn A on calcium current in ventricular cells of guinea pig. *Acta Pharmacol Sin* 1994; 15: 525-7.
- 8 Sun F, Li DX. Effect of *l*-tetrahydropalmatine on isolated guinea pig atria. *Chin Pharmacol Bull* 1989; 5: 158-61.

329-331

四氢巴马汀对豚鼠心室肌细胞跨膜内向  $Ca^{2+}$  电流

R 965.1 R 972

## 的抑制作用

徐春, 孙明智, 李玉荣, 杨宝峰<sup>1</sup>, 王丽娟<sup>2</sup>, 李金鸣<sup>2</sup>(哈尔滨医科大学生理教研室, <sup>1</sup> 药理教研室, 哈尔滨 150086; <sup>2</sup> 中国医科大学药理教研室, 沈阳 110001, 中国)

关键词 四氢巴马汀; 心肌; 钙通道; 膜片钳技术; 培养的细胞培养

**A 目的:** 研究四氢巴马汀 (THP) 对豚鼠心室肌细胞膜  $Ca^{2+}$  通道的作用 **方法:** 分离豚鼠心室肌细胞, 利用膜片钳技术观察、记录心室肌细胞膜跨膜内向钙电流 ( $I_{Ca}$ ) **结果:** THP 灌流 5 min 后明显抑制豚鼠心室肌细胞内向  $I_{Ca}$ , 其作用呈现剂量-效应关系, 并具有频率依赖性. 0.1, 1, 10  $\mu\text{mol} \cdot \text{L}^{-1}$  的 THP 使  $I_{Ca}$  从  $1.15 \pm 0.22$ ,  $0.91 \pm 0.18$ ,  $1.60 \pm 0.42$  nA (control,  $n=5$ ) 分别降低到  $0.90 \pm 0.21$  ( $P < 0.01$ ),  $0.56 \pm 0.21$  ( $P < 0.01$ ),  $0.83 \pm 0.21$  ( $P < 0.05$ ) nA, 抑制率分别为 22%, 38% 和 48%, THP 的作用易于洗脱. 峰值电压为 0 mV, 此时 THP 作用最明显, THP 对较高频率刺激产生的电流抑制作用较强, 2 Hz 时的抑制率高于 0.1 Hz 时的抑制率 ( $P < 0.01$ ) **结论:** THP 具有  $Ca^{2+}$  通道阻滞作用.

## Effect of sodium glycyrrhetinate on neonatal rat myocardial cells

ZHU Xiao-Wei, MENG Fu-Ming

(Department of Pharmacology, Lanzhou Medical College, Lanzhou 730000, China)

**KEY WORDS** glycyrrhetinate; myocardium; cultured cells; cyclic AMP; calcium; oxygen

**AIM:** To investigate the effect of sodium glycyrrhetinate (SG) on neonatal rat myocardial cells. **METHODS:** The neonatal rat myocardial cells were cultured *in vitro*. Radioimmunoassay and fluorimetry were used to determine cAMP and  $[Ca^{2+}]$ , respectively. **RESULTS:** The beating

rate of myocardial cells was depressed by SG 0.4  $\text{mmol} \cdot \text{L}^{-1}$  at 5, 10, and 15 min, from  $73 \pm 9 \text{ min}^{-1}$  to  $62 \pm 5$ ,  $59 \pm 7$ , and  $56 \pm 6 \text{ min}^{-1}$ , respectively. SG 0.1 and 0.2  $\text{mmol} \cdot \text{L}^{-1}$  showed above similar results at 15 min and 10, 15 min, respectively. When the myocardial cells were incubated with SG 0.2 and 0.4  $\text{mmol} \cdot \text{L}^{-1}$  at 37 °C for 10 and 15 min, the concentration of cAMP and  $[Ca^{2+}]$  were reduced. cAMP contents in SG 0.2  $\text{mmol} \cdot \text{L}^{-1}$  treated group at 10 and 15 min were lower than

control group ( $1.09 \pm 0.18$  vs  $1.65 \pm 0.48$  pmol per vial,  $P < 0.05$ ;  $1.12 \pm 0.35$  vs  $1.72 \pm 0.49$  pmol per vial,  $P < 0.01$ ), and so was  $[Ca^{2+}]$  ( $30 \pm 4$  nmol·L<sup>-1</sup> vs  $41 \pm 6$  nmol·L<sup>-1</sup>,  $P < 0.05$  and  $28 \pm 6$  nmol·L<sup>-1</sup> vs  $38 \pm 7$  nmol·L<sup>-1</sup>,  $P < 0.01$ ). SG 0.1 and 0.2 nmol·L<sup>-1</sup> increased the pO<sub>2</sub> change rate ( $87\% \pm 5\%$ ,  $75\% \pm 4\%$  vs  $54\% \pm 3\%$ ,  $P < 0.01$ ) in suspension fluid of myocardial cells, but SG 0.4 mmol·L<sup>-1</sup> decreased it ( $31\% \pm 2\%$  vs  $54\% \pm 3\%$ ,  $P < 0.01$ ). **CONCLUSION:** SG protects myocardium or treats ischemic cardiac disease.

Glycyrrhetinate, the hydrolysate of glycyrrhizine, had pharmacological actions of anti-inflammation, scavenging oxygen free radicals, and immunomodulating function<sup>[1,2]</sup>. Sodium glycyrrhetinate (SG) decreased the heart rate of anesthetized rat<sup>[3]</sup>. In this experiment, its effects on myocardial cells were observed.

**MATERIALS**

SG (colorless powder, purity: 98.5%), made by Ningxia Institute of Chemical Industry, was dissolved in 10% Me<sub>2</sub>SO (Sigma Chemical Co). Medium DMEM was made by Gibco. The cyclic [<sup>3</sup>H]AMP box was purchased from Institute of Basic Medical Sciences, Chinese Academy of Medical Sciences. Fura 2-AM (light yellow powder, AR grade), synthesized by Institute of Materia Medica, Chinese Academy of Medical Sciences, was dissolved in Me<sub>2</sub>SO. Wistar rats were inbred in Research Centre of Experimental Animals, Chinese Academy of Medical Sciences, and were fed to bear neonatal rat.

**METHODS AND RESULTS**

**Myocardial cell cultures**<sup>[4]</sup> Heart ventricles from 2 - 5 d old rat were digested with 0.06%

trypsin. The suspension of myocardial cells was purified<sup>[5]</sup>. Myocardial cells were dispersed and cultured on DMEM containing 15% fetal bovine serum at 37 °C with 95% O<sub>2</sub> + 5% CO<sub>2</sub>.

**Effect of SG on beating rate** After myocardial cells were cultured under standard condition for 5 - 10 d, the culture flasks which contained SG 0.1, 0.2, and 0.4 mmol·L<sup>-1</sup> and 10% Me<sub>2</sub>SO 2 mL·L<sup>-1</sup> (all of final concentrations) were placed under a microscope. At  $37.0 \pm 0.2$  °C, SG 0.4 mmol·L<sup>-1</sup> decreased the beating rates at 5, 10, and 15 min. SG 0.1 and 0.2 mmol·L<sup>-1</sup> showed similar effects at 15 min and 10, 15 min, respectively (Tab 1).

**Effect of SG on cAMP** Having been cultured for 5 - 10 d, myocardial cells were incubated with SG and 10% Me<sub>2</sub>SO at 37 °C for 10 and 15 min. The medium was thrown away and cAMP was extracted with HClO<sub>4</sub><sup>[6]</sup> and measured by radioimmunoassay. SG 0.2 and 0.4 mmol·L<sup>-1</sup> decreased the cAMP concentration in myocardial cells ( $P < 0.05$ ) (Tab 1).

**Effect of SG on  $[Ca^{2+}]$**  The myocardial cells prepared were loaded with Fura 2-AM<sup>[7]</sup>, and were treated with SG and 10% Me<sub>2</sub>SO at 37 °C for 10 or 15 min. The Ca<sup>2+</sup> concentration was determined fluorimetrically<sup>[8]</sup>. It was shown that SG 0.2 and 0.4 mmol·L<sup>-1</sup> reduced the intracellular free Ca<sup>2+</sup> concentration ( $P < 0.01$  or 0.05). (Tab 1)

**Effect of SG on pO<sub>2</sub>**<sup>[9]</sup> SG 0.1 and 0.2 mmol·L<sup>-1</sup> increased the rate of pO<sub>2</sub> changing ( $P < 0.01$ ), but SG 0.4 mmol·L<sup>-1</sup> decreased it ( $P < 0.01$ ). (Tab 1)

**DISCUSSION**

SG concentration-dependently decreased the

**Tab 1. Effects of sodium glycyrrhetinate on neonatal rat myocardial cells.** n = numbers of culture flasks.  $\bar{x} \pm s$ . \* $P > 0.05$ , <sup>b</sup> $P < 0.05$ , <sup>c</sup> $P < 0.01$  vs control.

SG/ mmol·L <sup>-1</sup>	0	Number of cell beats per minute					cAMP/pmol per vial		$[Ca^{2+}]$ /nmol·L <sup>-1</sup>		pO <sub>2</sub> change/%
		1	3	5	10	15	10	15	10	15 min	
0	71±7 (6)	73±7	70±11	72±7	73±8	74±10	1.65±0.48 (6)	1.72±0.49 (5)	41±6 (5)	38±7 (5)	54±3 (5)
0.4	73±9 (5) <sup>a</sup>	74±7 <sup>a</sup>	68±9 <sup>a</sup>	62±6 <sup>b</sup>	59±7 <sup>c</sup>	56±6 <sup>c</sup>	0.92±0.32 (5) <sup>b</sup>	1.08±0.28 (5) <sup>b</sup>	20±4 (5) <sup>c</sup>	22±6 (5) <sup>c</sup>	31±2 (6) <sup>c</sup>
0.2	74±6 (5) <sup>a</sup>	71±7 <sup>a</sup>	70±6 <sup>a</sup>	67±6 <sup>a</sup>	62±3 <sup>b</sup>	60±4 <sup>b</sup>	1.09±0.18 (5) <sup>b</sup>	1.12±0.35 (5) <sup>b</sup>	30±4 (5) <sup>c</sup>	28±6 (5) <sup>b</sup>	75±4 (5) <sup>c</sup>
0.1	75±9 (5) <sup>a</sup>	73±10 <sup>a</sup>	71±4 <sup>a</sup>	68±4 <sup>a</sup>	66±7 <sup>a</sup>	61±5 <sup>b</sup>	1.29±0.42 (5) <sup>a</sup>	1.26±0.41 (5) <sup>b</sup>	36±5 (5) <sup>a</sup>	30±7 (5) <sup>a</sup>	87±5 (5) <sup>c</sup>

beating rate of myocardial cells, which is related to the decrease of intracellular cAMP and  $Ca^{2+}$  concentration<sup>(10)</sup>. Influence of SG in high or low concentration on the oxygen consumption of myocardial cells was different. It suggests that SG in low concentration improve the catabolic metabolism, like its structural amagluocortid<sup>(11)</sup>. The effect of SG on decreasing the rate of  $pO_2$  changing in high concentration may be related to the sharp decrease of beating rate. Above all, it is considered that SG may be used to protect myocardium or treat ischemic cardiac disease.

## REFERENCES

- 1 Li XF, Meng FM, Pan FS, Lun W, Lan ZF, Wang FL, *et al.* Effects of 18 $\beta$ -sodium glycyrrhetic acid (SGA) on inflammation and immune function. *Chin Pharmacol Bull* 1990; 6: 105-8.
- 2 Ju HS, Li XJ, Zhao BL, Hou JW, Han ZW, Xin WJ. Scavenging effects of sodium ferulate and 18 $\beta$ -glycyrrhetic acid on oxygen free radicals. *Acta Pharmacol Sin* 1990; 11: 466-70.
- 3 Li XF, Wu YJ, Guo ZH, Liu L. Effect of sodium 18 $\beta$ -glycyrrhetate on experimental arrhythmia. *Chin J Chin Mat Med* 1992; 17: 176-8.
- 4 Liu JJ, Yang YC, Zhu QY, Gu KM, Tan SJ. An improvement on the culturing method of myocardial cells. *Acad J Sec Mil Med Univ* 1987; 8: 367-8.
- 5 Sun H. The purifying method of cultured myocardial cells *in vitro* and its application. *Chin J Cell Biol* 1988; 10: 85-7.
- 6 Miao CY, Gu KM. Effects of sulmazole on beating rate and cAMP levels in cultured cardiomyocytes. *Acta Pharmacol Sin* 1990; 11: 217-20.
- 7 Mier WG, Cannell MB, Berlin JR, Marban E, Lederer WJ. Cellular and subcellular heterogeneity of  $[Ca^{2+}]$  in single heart cells revealed by Fura-2. *Science* 1987; 235: 325-7.
- 8 Pan JY, Huang XD, Zhan CY. Measurement of the intracellular free calcium concentration of ventricular myocytes with fluorescent dye Fura-2. *J Sun Yat-Sen Med Univ* 1990; 11: 14-6.
- 9 Li LD, Li YO, Sun H, Gao FH, Zhang JM, Ma YL, *et al.*

Influences of 34 chinese herbal medicines promoting blood circulation to remove blood stasis on the oxygen consumption of cultured neonatal myocardial cells.

*Chin J Inter Trad West Med* 1989; 9: 292

- 10 Drummond GI, Severson DL. Cyclic nucleotides and cardiac function. *Circ Res* 1979; 44: 145-53.
- 11 White DA, Middleton B, Baxter M. Hormones and metabolic control. A medical student's guide to control of various aspects of normal and abnormal metabolism. London: Edward Arnold, 1984: 1-16

31-333

甘草次酸钠对乳鼠心肌细胞的影响

R=85.5

R=82.710

朱晓卫, 孟富敏

(兰州医学院药理学教研室, 兰州 730000, 中国)

关键词 甘草次酸; 心肌; 培养的细胞; 腺苷环磷(酸); 钙; 氧 细胞培养, 甘草

目的: 研究甘草次酸钠(SG)对乳鼠心肌细胞的作用。方法: 体外培养乳鼠心肌细胞, 用放射免疫法和荧光法分别测定 cAMP 和  $[Ca^{2+}]$ , 结果: SG  $0.4 \text{ mmol} \cdot \text{L}^{-1}$  于 5, 10 和 15 min 使心肌细胞搏动频率由  $73 \pm 9 \text{ min}^{-1}$  分别降至  $62 \pm 5$ ,  $59 \pm 7$  和  $56 \pm 6 \text{ min}^{-1}$ 。SG  $0.1$  和  $0.2 \text{ mmol} \cdot \text{L}^{-1}$  分别在 15 和 10, 15 min 呈现以上相似的结果。SG  $0.2$  和  $0.4 \text{ mmol} \cdot \text{L}^{-1}$  与心肌细胞  $37^\circ\text{C}$  共孵 10, 15 min, cAMP 浓度和  $[Ca^{2+}]$  均被降低。SG  $0.2 \text{ mmol} \cdot \text{L}^{-1}$  处理组的 cAMP 含量低于对照组 ( $1.09 \pm 0.18$ ,  $1.12 \pm 0.35 \text{ pmol per vial}$ ,  $P < 0.05$ );  $[Ca^{2+}]$  的变化也类似 ( $30 \pm 4 \text{ nmol} \cdot \text{L}^{-1}$   $P < 0.01$ ,  $28 \pm 6 \text{ nmol} \cdot \text{L}^{-1}$   $P < 0.05$ )。SG  $0.1$  和  $0.2 \text{ mmol} \cdot \text{L}^{-1}$  提高心肌细胞悬液氧分压变化率 ( $87\% \pm 5\%$ ,  $75\% \pm 4\%$ ,  $P < 0.01$ ), 但 SG  $0.4 \text{ mmol} \cdot \text{L}^{-1}$  降低氧分压变化率 ( $31\% \pm 2\%$ ,  $P < 0.01$ )。结论: SG 可保护心肌或治疗缺血性心脏病。