

Pinacidil suppression on 5-HT₃ receptor-mediated contraction of guinea pig ileum *in vitro*

YUAN Yu-Min, XU De-Yi¹, HU Guo-Yuan² (State Key Laboratory of Drug Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 200031, China)

KEY WORDS serotonin receptors; pinacidil; ileum; myenteric plexus; carbachol; GR65630

AIM: To study the effects of the K⁺ channel opener pinacidil on 5-HT₃ receptor-mediated contractions of the isolated guinea pig ileum (GPI) longitudinal muscle-myenteric plexus strip preparations.

METHODS: GPI contractions were recorded with a chart recorder through isometric transducers. The effect of pinacidil on binding properties of 5-HT₃ receptors was assessed using [³H]GR65630 binding assay in membrane preparations of rat entorhinal cortex. **RESULTS:** (1) A selective 5-HT₃ receptor agonist 2-methyl-5-HT 0.1–300 μmol·L⁻¹ and 5-HT 0.001–50 μmol·L⁻¹ elicited GPI contractile responses in concentration-dependent manners, the EC₅₀ values (and 95 % confidence limits) for 2-methyl-5-HT and 5-HT were 10.0 (8.9–11.2) μmol·L⁻¹ and 1.6 (1.3–1.9) μmol·L⁻¹, respectively. Selective 5-HT₃ receptor antagonist tropisetron 0.1 μmol·L⁻¹ competitively inhibited the responses to 2-methyl-5-HT and 5-HT. (2) Pinacidil 0.5–5 μmol·L⁻¹ inhibited 5-HT₃ receptor-mediated contractions. (3) Pinacidil 1 μmol·L⁻¹ enhanced the inhibitory effects of tropisetron 0.1 μmol·L⁻¹ or another selective 5-HT₃ receptor antagonist benesetron 1 μmol·L⁻¹ on 5-HT-induced GPI contractile responses. (4) Pinacidil 1–5 μmol·L⁻¹ did not affect GPI contractile responses evoked by a selective M-ACh receptor agonist carbachol 1 μmol·L⁻¹. (5) Pinacidil 1–5 μmol·L⁻¹ had no effect on binding properties of 5-HT₃ receptors with selective 5-HT₃ receptor radioligand [³H]GR65630 in the entorhinal cortex of rat brain. **CONCLUSION:** The inhibition by pinacidil of 5-HT₃ receptor-mediated GPI contractile responses may be mediated through activation of ATP-

sensitive K⁺ channels located in prejunctional myenteric neurons.

The K⁺ channel openers (KCO) form a novel class of compounds in clinical applications including the treatment of cardiovascular diseases, bronchial asthma, or irritable bladder syndrome. These KCO acted primarily on a heterogeneous class of K⁺ channels, ATP-sensitive K⁺ channels, which were found in excitable tissues including heart, skeletal and smooth muscle, neurons, and pancreatic β-cells^[1]. Basically, these agents increase potassium conductance by activation of ATP-sensitive K⁺ channels and thus lead to a hyperpolarization of the cell membrane and reduced membrane excitability. Consequently, KCO relax smooth muscle preparations precontracted by spasmogens^[2].

5-HT₃ receptor was originally described as 'M' type of 5-HT receptor in view of the inhibition by morphine of the contractile response of isolated GPI to 5-HT. In electrophysiologic studies, 5-HT₃ receptors have been characterized as a new member of the ligand-gated ion channel superfamily, similar in many aspects to N-ACh receptor^[3]. The function of 5-HT₃ receptors in both the central and peripheral nervous systems is mainly involved in modulation of the release of neurotransmitters, including ACh, dopamine, GABA, noradrenaline, cholecystokinin, etc^[4]. The main component of GPI contractions induced by 5-HT 0.1–10 μmol·L⁻¹ was sensitive to tetrodotoxin 1 μmol·L⁻¹, thus mediated via activation of neuronal 5-HT₃ receptors that led to the release of ACh from the myenteric plexus neurons^[5,6].

Cromakalim, a K⁺ channel opener, reduced cholinergic transmission in airway smooth muscle preparations^[7] and inhibited contractions of GPI caused by the release of endogenous ACh via electric stimulation or via activation of 5-HT₃ receptors^[8,9]. Pinacidil, a cyanoguanidine potassium channel opener, inhibited release of dopamine and norepinephrine in rat

¹ Visiting researcher from Nanjing Railway Medical College.

² Correspondence to Prof HU Guo-Yuan.

Phn 86-21-6471-8166. Fax 86-21-6473-3473.

Received 1997-04-07

Accepted 1997-07-02

vas deferens¹⁰. In the present work, the effects of pinacidil on the 5-HT₃ receptor-mediated contractile responses of GPI were examined.

MATERIALS AND METHODS

Chemicals 2-Methyl-5-HT, benesetron and pinacidil (Research Biochemicals International, USA); 5-HT, carbachol and metoclopramide (Sigma, USA); [³H]GR65630 (DuPont NEN, USA); tropisetron (synthesized and gifted by Prof ZHU You-Cheng, Shanghai Institute of Materia Medica, Chinese Academy of Sciences).

Measurement of contractions of GPI Guinea pigs (♂, 300–400 g) were stunned. The ileum was excised approximately 10 cm from the ileo-cecal junction and strips of longitudinal muscle with adherent myenteric plexus were freed from the underlying circular muscle. In each organ bath (5 mL), one longitudinal muscle strip, 2–3 cm long, was mounted. The bath contained Tyrode's solution, which consisted of: NaCl 137.0, CaCl₂ 1.8, KCl 2.7, MgCl₂ 1.05, NaHCO₃ 11.9, NaH₂PO₄ 0.4, glucose 5.6 mmol·L⁻¹, and gassed with 95% O₂ + 5% CO₂ at 37 °C. After equilibration for 1 h, the strips were placed under a resting tension of 0.5 g. Contractions were recorded with a chart recorder via isometric transducers. Noncumulative concentration-response curves were established by adding increasing concentrations of 5-HT₃ receptor agonist (50 μL) to the organ bath at intervals of at least 20 min, which was found in preliminary experiments to be long enough to avoid tachyphylaxis. Each concentration of the agonist was left in contact with the tissue for about 30 s. The curves were then reconstructed in the presence of pinacidil with increasing concentrations applied 5 min before each application of the agonist. Each strip was used to record only 2 concentration-response curves: one for the agonist alone and the other for the agonist in the presence of one concentration of pinacidil. Experiments with the same protocol were repeated with selective 5-HT₃ receptor antagonists tropisetron or benesetron added 10 min before agonist application. All peak responses to each concentration of the agonist were expressed as a % of the response to carbachol 10 μmol·L⁻¹.

Receptor-binding assay with radioligand Sprague-Dawley rats (♂, 200–250 g) were decapitated. Pooled entorhinal cortex tissue was homogenized (Ultra Turrax T25) in 10 vol HEPES buffer (50 mmol·L⁻¹, pH 7.4, 4 °C) and spun at 4 °C and 49 000 × g for 10 min. The supernatant was discarded and the process repeated for the pellet. The final pellet was suspended in 10 vol HEPES buffer. For binding, assay tubes were added with 100 μL selective 5-HT₃ receptor radioligand [³H]GR65630 (2286.6 TBq·mol⁻¹, final concentration of 0.3 nmol·L⁻¹) in HEPES buffer, testing drug 100 μL, and tissue preparation 100 μL (0.3–0.4 mg protein). Tubes were incubated at 37 °C for 30 min, which was terminated

by rapid vacuum filtration through Whatman GF/B filters using a Brandel Cell Harvester. Filters were washed immediately with 3 mL Tris·HCl buffer (50 mmol·L⁻¹, pH 7.4, 4 °C) for three times. Radioactivity was assayed with a Beckman LS6000LL scintillometer (efficiency = 47%). All individual assays were carried out in replicates of three. Nonspecific binding was determined by the inclusion of 5-HT₃ receptor antagonist metoclopramide 100 μmol·L⁻¹, which inhibited 50%–60% of total binding of [³H]GR65630 0.3 nmol·L⁻¹ in washed crude homogenates of rat entorhinal cortex.

Data analysis EC₅₀ and IC₅₀ values were calculated using computer software 'GraphPad InPlot'. The data were presented as $\bar{x} \pm s$ and compared with *t*-test.

RESULTS

5-HT₃ receptor-mediated GPI contraction

2-Methyl-5-HT, a selective 5-HT₃ receptor agonist, evoked GPI contractile responses in a concentration-dependent manner with EC₅₀ of 10.0 (8.9–11.2) μmol·L⁻¹. Selective 5-HT₃ receptor antagonist tropisetron 0.1 μmol·L⁻¹ competitively inhibited 2-methyl-5-HT induced contractile responses. Similar phenomena were seen when 5-HT was used as agonist (Fig 1). The EC₅₀ for 5-HT was 1.6 (1.3–1.9) μmol·L⁻¹. Note that the responses induced by 5-HT (1–10 nmol·L⁻¹) were insensitive to tropisetron. Only the responses induced by 5-HT (≥ 0.1 μmol·L⁻¹) were mediated via 5-HT₃ receptors.




Fig 1. 2-Methyl-5-HT- and 5-HT-induced GPI contractions with and without tropisetron 0.1 μmol·L⁻¹. *n* = 5 strips, $\bar{x} \pm s$.

Effects of pinacidil on 5-HT₃ receptor-mediated GPI contraction Pinacidil inhibited GPI

contractile responses to 2-methyl-5-HT in a concentration-dependent manner. Increasing concentrations of pinacidil from 0.5 to 5 $\mu\text{mol}\cdot\text{L}^{-1}$ produced progressive suppression of GPI contraction induced by 2-methyl-5-HT. Addition of pinacidil into the organ bath always caused a substantial decrease of spontaneous activity of GPI preparations (Fig 2). Similar results were obtained when 5-HT 0.1 – 10 $\mu\text{mol}\cdot\text{L}^{-1}$ was used as agonist.




Fig 2. Suppression of 2-methyl-5-HT-induced GPI-contraction by pinacidil. $n = 5$ strips, $x \pm s$. Concurrent contractions are shown for 2-methyl-5-HT 10 $\mu\text{mol}\cdot\text{L}^{-1}$ with and without pinacidil 1 $\mu\text{mol}\cdot\text{L}^{-1}$.

Pinacidil enhanced the inhibitory effect of tropisetron and benesetron on 5-HT-evoked GPI contraction GPI contractions induced by 5-HT (0.1 – 10 $\mu\text{mol}\cdot\text{L}^{-1}$) were competitively suppressed by tropisetron 0.1 $\mu\text{mol}\cdot\text{L}^{-1}$. When pinacidil 1 $\mu\text{mol}\cdot\text{L}^{-1}$ was added, the inhibitory effect of tropisetron increased. Similar results were obtained when another selective 5-HT₃ receptor antagonist benesetron 1 $\mu\text{mol}\cdot\text{L}^{-1}$ was tested (Fig 3).

Effect of pinacidil on carbachol-evoked GPI contraction Pinacidil 1 and 5 $\mu\text{mol}\cdot\text{L}^{-1}$, which suppressed GPI contractions induced by 2-methyl-5-HT (Fig 2), had no detectable effects on GPI contractile responses induced by a selective M-ACh receptor agonist carbachol 1 $\mu\text{mol}\cdot\text{L}^{-1}$. The carbachol-induced contractions in presence of pinacidil 1 or 5 $\mu\text{mol}\cdot\text{L}^{-1}$ (68% \pm 14%, 71% \pm 12%) had no significant difference with those induced by carbachol alone (74% \pm 12%). These observations suggest that the inhibition by pinacidil of 5-HT₃ receptor-mediated GPI contractions may be mediated via a prejunctional mechanism.




Fig 3. Enhancement of inhibitory effects of tropisetron 0.1 $\mu\text{mol}\cdot\text{L}^{-1}$ or benesetron 1 $\mu\text{mol}\cdot\text{L}^{-1}$ on 5-HT-induced contraction of GPI by pinacidil 1 $\mu\text{mol}\cdot\text{L}^{-1}$. $n = 3$ strips, $x \pm s$. $^aP > 0.05$, $^bP < 0.05$, $^cP < 0.01$ vs inhibition by tropisetron or benesetron alone.

Effect of pinacidil on binding properties of 5-HT₃ receptors Due to the relatively low density of 5-HT₃ receptors in myenteric plexus neurons of GPI, we tested the effect of pinacidil on the binding properties of 5-HT₃ receptors in rat entorhinal cortex that has been found to have the highest distribution of 5-HT₃ receptors in rat brain^[11]. Pinacidil 5 $\mu\text{mol}\cdot\text{L}^{-1}$ neither inhibited the total binding of 5-HT₃ receptor radioligand [³H]GR65630 0.3 $\text{nmol}\cdot\text{L}^{-1}$, nor affect the IC₅₀ of 5-HT, 2-methyl-5-HT, tropisetron, and benesetron to compete for [³H]GR65630 0.3 $\text{nmol}\cdot\text{L}^{-1}$ binding (Tab 1). These results suggested that pinacidil had no direct effect on the property of 5-HT₃ receptor binding for [³H]GR65630.

Tab 1. Effects of pinacidil on [³H] GR65630 0.3 $\text{nmol}\cdot\text{L}^{-1}$ binding with 5-HT₃ receptors. $n = 3$ membrane homogenates of 5 rat entorhinal cortices, $x \pm s$. $^aP > 0.05$ vs control.

	Control	Pinacidil (5 $\mu\text{mol}\cdot\text{L}^{-1}$)
Total binding/Bq	71 \pm 6	69 \pm 7 ^a
IC ₅₀ / $\text{nmol}\cdot\text{L}^{-1}$		
5-HT	208 \pm 45	228 \pm 56 ^a
2-Methyl-5-HT	156 \pm 37	166 \pm 35 ^a
Tropisetron	18 \pm 4	23 \pm 7 ^a
Benesetron	99 \pm 21	118 \pm 29 ^a

DISCUSSION

The results of the present study showed that

pinacidil within a certain range of concentrations inhibited the 5-HT₃ receptor-mediated GPI contractile responses. These effects seemed not to be produced via a postjunctional mechanism since the carbachol-induced GPI contractile responses were not affected by the K⁺ channel opener. The action site of pinacidil is likely to be in prejunctional myenteric plexus neurons where 5-HT₃ receptors locate as suggested by Zimi *et al* that the effect of KCO could occur presynaptically^[11]. This notion was also in accord with the previous report that 5-HT-induced GPI contractile responses induced by 5-HT 0.1 - 10 μmol·L⁻¹ were mediated via neuronally located 5-HT₃ receptors^[12]. On the other hand, Sun and Benishin found that pinacidil 1 μmol·L⁻¹ decreased the basal tension in the plexus-free longitudinal muscle of guinea-pig ileum^[13]. This observation has thus been implicated that the mechanisms of action of pinacidil might be involved in ATP-sensitive K⁺ channels on postjunctional smooth muscle cells. However, to date there has been no direct ligand-binding and electrophysiological evidence for the presence of ATP-sensitive K⁺ channels in the plexus-free longitudinal muscle cells of guinea pig ileum.

Our results showed that the effects of pinacidil on 5-HT₄-mediated contractions of GPI seemed not due to changes in the 5-HT₃ receptors *per se*, at least in their property of binding for the specific ligand. When 5-HT₄ receptors in myenteric plexus neurons were activated by the agonists, Na⁺ and K⁺ fluxes through the cation-selective channels coupled directly to the receptors caused a rapid depolarization that in turn activated voltage-gated Ca²⁺ channels and caused an increase in the cytosolic free Ca²⁺ concentration^[4], then led to neurotransmitter release which caused GPI contractions^[6]. Pinacidil may open the ATP-sensitive K⁺ channels in the same neurons, thus counteract the 5-HT₃ receptor-mediated effects. The fact that spontaneous activity of GPI greatly decreased after addition of pinacidil supported the above speculation. The enhancement by pinacidil of the inhibitory effect of tropisetron or benesetron on the 5-HT-evoked GPI contractile responses, therefore, could be attributed to a certain type of functional synergism. However, cromakalim, another K⁺ channel opener, did not affect the release of [³H]-acetylcholine evoked by electric stimulation or by activation of 5-HT₃ receptors

in GPI^[9]. In addition, there are recently some debates about the effects of KCO on Ca²⁺ entry which causes contractions of various smooth muscle preparations^[2]. The effects of KCO on intestinal muscle are still not fully understood.

Nevertheless, our results suggested that some functional interactions exist between 5-HT₃ receptors and ATP-sensitive K⁺ channels in longitudinal muscle-myenteric plexus neurons of GPI. The physiological significance of this finding needs to be addressed at single neuron level using new techniques, such as electrophysiological study, intracellular Ca²⁺ imaging, *etc.*

ACKNOWLEDGMENTS To Ms YANG Hui-Hua for her excellent technical assistance. This study was partly supported by a research grant from Chinese Academy of Sciences (KZ 951-A1-103).

REFERENCES

- 1 Edwards G, Weston AH. The pharmacology of ATP-sensitive potassium channels. *Annu Rev Pharmacol Toxicol* 1993; 33: 597-637.
- 2 Quast U. Do the K⁺ channel openers relax smooth muscle by opening K⁺ channels? *Trends Pharmacol Sci* 1993; 14: 332-7.
- 3 Derkach V, Surprenant A, North RA. 5-HT₃ receptors are membrane ion channels. *Nature* 1989; 339: 706-9.
- 4 Jackson MB, Yakel JL. The 5-HT₃ receptor channel. *Annu Rev Physiol* 1995; 57: 447-68.
- 5 Buchheit KH, Engel G, Mutschler E, Richardson B. Study of the contractile effect of 5-hydroxytryptamine (5-HT) in the isolated longitudinal muscle strip from guinea-pig ileum. *Naunyn-Schmiedeberg's Arch Pharmacol* 1985; 329: 36-41.
- 6 Fox AJ, Morton IKM. An examination of the 5-HT₃ receptor mediating contraction and evoked [³H]-acetylcholine release in the guinea-pig ileum. *Br J Pharmacol* 1990; 101: 553-6.
- 7 McCaig DJ, Jonckheere BD. Effect of cromakalim on bronchoconstriction evoked by cholinergic nerve stimulation in guinea-pig isolated trachea. *Br J Pharmacol* 1989; 93: 662-8.
- 8 Zimi S, Ben-Ari Y, Ashford MLJ. Characterization of sulfonyleurea receptors and the action of potassium channel openers on cholinergic neurotransmission in guinea pig isolated small intestine. *J Pharmacol Exp Ther* 1991; 259: 566-73.
- 9 Schwörer H, Kithinger H. Effects of cromakalim on acetylcholine release and smooth muscle contraction in guinea-pig small intestine. *Naunyn-Schmiedeberg's Arch Pharmacol* 1989; 339: 706-8.
- 10 Soares-da-Silva P, Fernandes MH. Inhibition by the putative potassium channel opener pinacidil of the electrically-evoked release of endogenous dopamine and noradrenaline in the rat vas deferens. *Naunyn-Schmiedeberg's Arch Pharmacol* 1990; 342: 115-21.
- 11 Kilpatrick GJ, Jones BJ, Tyers MB. Identification and distribution

of 5-HT₃ receptors in rat brain using radioligand binding.
Nature 1987; 330: 746-48

- 12 Eglén RM, Swank SR, Walsh LKM, Whiting RL.
Characterization of 5-HT₃ and 'atypical' 5-HT receptors mediating
guinea-pig ileal contractions *in vitro*.
Br J Pharmacol 1990; 101: 513-20.
- 13 Sun YD, Benishin CG. K⁺ channel openers relax longitudinal
muscle of guinea pig ileum. Eur J Pharmacol 1994; 271: 453-9.

31-35

吡那地尔对 5-HT₃ 受体介导的 离体豚鼠回肠收缩反应的抑制

袁玉民¹, 许德义¹, 胡国渊² (中国科学院上海药物
研究所新药研究国家重点实验室, 上海 200031, 中国)

关键词 血清素受体; 吡那地尔; 回肠; 肠肌丛;
卡巴胆碱; GR65630

R972
R975-6

收缩反应

目的: 研究钾通道开放剂吡那地尔对 5-HT₃ 受体介导的离体豚鼠回肠(GPI)收缩反应的影响。方法: 以等长换能器记录 GPI 收缩反应, 以 [³H]GR65630 结合试验检测大鼠内嗅皮层 5-HT₃ 受体的结合特性。结果: (1) 2-甲基-5-HT 及 5-HT 以剂量依赖方式引起 GPI 的收缩; 托品色创竞争性抑制此反应。(2) 吡那地尔以剂量依赖方式抑制 2-甲基-5-HT 和 5-HT 引起的 GPI 收缩, 并增强托品色创或 Benesetron 对 5-HT 诱发 GPI 收缩的抑制作用, 但不影响卡巴胆碱引起的 GPI 收缩; 吡那地尔对大鼠内嗅皮层 5-HT₃ 受体与 [³H]GR65630 的结合无影响。结论: 吡那地尔可能通过激活突触前神经元 ATP 敏感钾通道抑制由 5-HT₃ 受体介导的 GPI 收缩反应。

Effects of long-term application of dopamine HCl on dopamine agonist-induced cAMP production in rat renal cortex¹

ZHAO Rong-Rui², JIN Xiao-Hong, ZHU Lin, WANG Wen-Ze
(Department of Physiology, Shanxi Medical University, Taiyuan 030001, China)

KEY WORD dopamine receptors; kidney cortex;
dopamine; cyclic AMP; dopamine agents; adenylyl
cyclase; Sch-23390; domperidone

AIM: To study the effects of long-term application of dopamine HCl (DA) on the functional changes of dopamine receptor subtypes coupled to adenylyl cyclase in rat renal cortex. **METHODS:** cAMP levels were measured by radioimmunoassay as an index of dopamine receptor function. **RESULTS:** Injection of DA (30 mg · kg⁻¹ · d⁻¹, ip, 30 d) reduced the fenoldopam (Fen) (100 μmol · L⁻¹)-induced increments of cAMP production from the control group of +1.26 ± 0.04 to the DA-treated group of +0.63 ± 0.22 nmol · min⁻¹/g tissue and the propyl-butyl-

dopamine (PBDA) (100 μmol · L⁻¹)-induced decrements of cAMP production in the presence of Sch-23390 (Sch) from the control group of -0.38 ± 0.18 to the DA-treated group of -0.11 ± 0.08 nmol · min⁻¹/g tissue with, however, comparable percentile changes for the 2 groups. Sch blocked both Fen- and PBDA-induced increase in cAMP production, while domperidone (Dom) blocked the decreasing effects of PBDA on cAMP accumulation in the presence of Sch. **CONCLUSION:** Long-term application of DA produced a marked "down regulation" of both DA₁ and DA₂ receptors in rat renal cortex with, however, the responsiveness of the remaining receptors unchanged.

Dopamine HCl (DA) agonists have been extensively used in the treatment of cardiovascular and kidney disease^[1,2]. But, long-term application of the drugs may result in the decrease of drug effects in patients and animals, indicating the occurrence of some

¹ Project supported by the National Natural Science Foundation of China, No. 81230150

² Correspondence to Prof ZHAO Rong-Rui. P. 036-351-413-5319.
Fax 036-351-202-1239

Received 1998-04-01

Accepted 1997-07-02