

Suppression of oxygen toxicity by melatonin

Russel J REITER¹, TAN Dun-Xian, QI Wen-Bo

(Department of Cellular and Structural Biology, The University of Texas Health Science Center, 7703 Floyd Curl Drive, San Antonio TX 78284-7762, USA)




KEY WORDS: melatonin; free radicals; antioxidants; oxidative stress; aging

ABSTRACT Melatonin, the chief secretory product of the pineal gland, was recently found to be a free radical scavenger and antioxidant. While most studies to date have used pharmacological quantities of melatonin to limit oxidative damage, physiologic concentrations of the indole which are present in aerobic organisms have also been shown to resist molecular damage inflicted by free radicals. Melatonin has several functions in terms of its antioxidative ability. It readily scavenges the most highly toxic free radical, the hydroxyl radical, and it directly detoxifies the peroxynitrite anion, nitric oxide, singlet oxygen, and the peroxyl radical. Precisely how efficient melatonin is in neutralizing each of these toxic agents remains to be determined. Melatonin also may stimulate several antioxidative enzymes including superoxide dismutase, glutathione peroxidase, and glutathione reductase as well as inhibiting the pro-oxidative enzyme, nitric-oxide synthase. Finally, melatonin chelates transition metal ions and prevents the deterioration of cellular membranes. This combination of actions may all contribute to melatonin's ability to reduce oxidative damage. Melatonin is highly effective in reducing nuclear DNA damage and membrane lipid destruction due to toxic free radicals *in vivo*. These findings have implications for disease processes, eg, neurodegenerative and cardiovascular diseases, which involve free radicals and for aging itself, which also is believed to be related to accumulated oxidative damage.

INTRODUCTION

Oxygen (O_2) is toxic because of its high electron (e^-) affinity which causes it to degrade


into a number of reactive and damaging intermediates. The reduction of O_2 produces a number of products which are either free radicals or products that form free radicals as follows:



Additionally, O_2 can add to an already existing radical ($R\cdot$) to yield a peroxyl radical:



Furthermore, O_2 forms a high-energy form, ie, singlet oxygen (1O_2):



A major mediator of oxygen toxicity is O_2^- which is formed *in vivo*⁽¹⁾. A primary site of O_2^- generation is the mitochondrion. During oxidative phosphorylation a small percentage of the e^- leak onto O_2 . Since O_2 accepts one e^- at a time, the first product is O_2^- (equation 1). A family of enzymes, the superoxide dismutases (SOD), which are present in cells of all aerobic organisms, reduce O_2^- to H_2O_2 (equation 2). While H_2O_2 is not a radical and, unless in high concentrations within cells, it *per se* is generally considered relatively unreactive. However, H_2O_2 does have a long half-life and easily passes through cell membranes; in so doing it can spread oxidative damage throughout the cell or even between cells. Its secondary toxicity derives from the fact that when H_2O_2 comes in contact with a transition metal ion, eg, Fe^{2+} , Cu^{1+} , it catalytically forms the $\cdot OH$ (equation 3). $\cdot OH$ is generally considered to be the most damaging radical generated from oxygen.

1O_2 is also not a free radical but it is capable of damaging tissues. This reactive species is sometimes used in the treatment of cancer in pho-

¹ Correspondence to Prof Russel J REITER.


Phn 1-210-576-3859. Fax 1-210-567-6948. E-mail Reiter@uthscsa.edu

Received 1998-09-13

Accepted 1998-09-18

tosensitization (photodynamic therapy). This treatment involves the administration of certain porphyrin derivatives to cancer patients; these derivatives preferentially localize in the tumor cells which are then exposed to visible light to generate $^1\text{O}_2$ which in turn damage the microvasculature of the tumor thereby leading to cell death.

Besides its dismutation to H_2O_2 , O_2^- can also combine with the gaseous neurotransmitter nitric oxide ($\text{NO}\cdot$) to form the peroxynitrite anion (ONOO^-).



ONOO^- , another nonradical product, is also highly toxic and furthermore it can degrade into the $\cdot\text{OH}$ or a $\cdot\text{OH}$ -like radical^[2].



Combating reactive oxygen species and free radicals are the function of antioxidants. Antioxidants can be either direct free radical scavengers, enzymes that metabolize reactive species to nontoxic products (eg, SOD) or they may function in the prevention of free radical generation (eg, transition metal ion chelators).

The most recently discovered molecule with antioxidant properties is melatonin. Melatonin is produced in several organs but blood levels of the indole are derived primarily from the pineal body in the brain^[3]. Unlike most other antioxidants, melatonin easily crosses all morphophysiological barriers, eg, placenta and blood-brain-barrier, and distributes to all parts of the cell where it provides protection against free radicals^[4]. While melatonin is generally known for its high lipid solubility^[5], it is also quite soluble in aqueous media^[6]. This review briefly summarizes the antioxidant activities of melatonin.

Melatonin as a direct free radical scavenger

The discovery that melatonin is a free radical scavenger was revolutionary and opened an area of investigation that has been aggressively pursued. While the initial suggestion that melatonin may detoxify free radicals appeared in 1991^[7], it was not until 1993 when the first definitive evidence was provided^[8]. In this study the authors used electron spin resonance spectroscopy (ESR) to show that melatonin scavenged the most toxic of the radicals, namely,

the $\cdot\text{OH}$. In the cell-free system melatonin successfully competed with the spin trap DMPO (5, 5-dimethylpyrroline-*N*-oxide) thereby reducing the adduct formed by the spin trapping agent and the $\cdot\text{OH}$; $\cdot\text{OH}$ were generated by exposing H_2O_2 to 254 nm ultraviolet light^[8].

Since this discovery, melatonin's ability to neutralize the $\cdot\text{OH}$ has been repeatedly confirmed using ESR and other technologies. Many verified the high $\cdot\text{OH}$ scavenging activity of melatonin *in vitro*^[9-11]. Melatonin *in vivo* also detoxifies the $\cdot\text{OH}$ ^[12]. These workers used microdialysis to collect extracellular fluid from the brain of rats treated with salicylate (which forms an adduct with the $\cdot\text{OH}$) and successfully showed that melatonin, just as in the *in vitro* studies, scavenges the $\cdot\text{OH}$.

The $\text{ROO}\cdot$ is highly damaging and propagates the chain reaction of lipid peroxidation. Melatonin is a more effective scavenger of the $\text{ROO}\cdot$ than vitamin E (tocopherol)^[13]. This is a remarkable claim considering the great efficacy of vitamin E in detoxifying the $\text{ROO}\cdot$. While melatonin is very effective in reducing the peroxidation of lipids, whether it functions as a chain breaking antioxidant by scavenging the $\text{ROO}\cdot$ is still uncertain. We^[14] and others^[15] have tested melatonin's ability to scavenge the $\text{ROO}\cdot$ and the conclusion from these studies is that it is not particularly efficient in scavenging the $\text{ROO}\cdot$. As will be noted below, however, melatonin is highly effective in resisting the oxidative breakdown of membrane lipids, especially *in vivo*^[16].

The high energy form of O_2 , ie, $^1\text{O}_2$, is toxic within cells and can cause extensive damage. It is usually generated by treating an animal with a photosensitizing agent and then exposing specific tissues to high intensity light. This method was used to determine the ability of melatonin to quench $^1\text{O}_2$ ^[17]. While the evidence presented is indirect in that melatonin prevented lipid peroxidation in the brain of rats treated with a photosensitizer and then exposed to light, the findings are consistent with melatonin quenching $^1\text{O}_2$. This finding has yet to be confirmed.

ONOO^- is a highly toxic agent formed by the combination of O_2^- and $\text{NO}\cdot$. Besides its inherent toxicity, ONOO^- also degrades into


other toxic agents including perhaps the $\cdot\text{OH}$ ⁽²⁾. Melatonin may also scavenges ONOO^- ⁽¹⁸⁾ as well as greatly reducing its toxicity *in vivo*⁽¹⁹⁾. Considering the important role of ONOO^- in promoting tissue damage in severe inflammatory states, melatonin's potential as an anti-inflammatory agent deserves additional study.

In a recent preliminary report it was also shown that melatonin directly scavenges $\text{NO}\cdot$ ⁽²⁰⁾. $\text{NO}\cdot$ functions as an important gaseous neurotransmitter but it can also be highly toxic under certain circumstances, for example, when it is excessively released during ischemia-reperfusion injury. The observation that melatonin scavenges $\text{NO}\cdot$ could potentially explain in part melatonin's protection against ischemia-reperfusion injury⁽²¹⁾.

While melatonin appears to be a rather ubiquitous free radical scavenger, it seems to lack the ability to directly detoxify O_2^- . On the other hand, it scavenges hypochlorous acid (HOCl)⁽¹⁴⁾. Both O_2^- and HOCl are generated by activated monocytes and macrophages when they engulf and kill bacteria. Melatonin actually promotes O_2^- formation by activated human monocytes^(22,23).

The products formed when melatonin scavenges a free radical are only beginning to be uncovered. Melatonin may work via e^- donation during which the indolyl cation radical is formed⁽²⁴⁾. This radical product was then proposed to combine with O_2^- to produce N^1 -acetyl- N^2 -formyl-5-methoxykynuramine.

Whereas the possibility has not been discounted that melatonin may function via e^- donation in the scavenging of the $\cdot\text{OH}$, three theoretical products may result from an alternative interaction⁽²⁵⁾. With the use of proton nuclear magnetic resonance ($^1\text{H-NMR}$), COSY $^1\text{H-NMR}$, mass spectrometry, and thermodynamic stability analysis we have identified one product of the interaction of melatonin with the $\cdot\text{OH}$ to be cyclic 3-hydroxymelatonin (unpublished observations by DX Tan, LC Manchester, RJ Reiter).



This product results after melatonin scavenges two $\cdot\text{OH}$ and we have shown that it appears in the urine of humans and other mammals.

Furthermore, its urinary concentrations are increased when animals are exposed to ionizing radiation which is known to generate the $\cdot\text{OH}$.

Melatonin's actions on anti- and pro-oxidative enzymes

A variety of enzymes play vital functions in the removal of reactive oxygen species from cells. As noted above, a family of SOD plays a central role in converting O_2^- to H_2O_2 , a nontoxic product. mRNA levels are stimulated following the exogenous administration of melatonin suggesting a possible rise in activity of SOD, although this was not actually documented⁽²⁶⁾.

The over expression of SOD without a commensurate rise in the H_2O_2 detoxifying enzyme, glutathione peroxidase (GPx), however, increases oxidative damage as seen in Down syndrome, amyotrophic lateral sclerosis, *etc.* Thus, the observations that melatonin also stimulates both mRNA for and the activity of GPx⁽²⁷⁾ was important since it assured the removal of the excess H_2O_2 that would result from the over stimulation of SOD. Furthermore, GPx also functions as an ONOO^- reductase⁽²⁸⁾, thereby removing this toxic agent from cells as well.

In the process of metabolizing H_2O_2 to nontoxic products, GPx oxidizes glutathione (GSH) to its disulfide form (GSSG) (Fig 1). So that GSH is not depleted from cells, GSSG is reduced to GSH in a reaction catalyzed by glutathione reductase (GRd). This replenishes GSH and allows for further metabolism of H_2O_2 (and other hydroperoxides) by GPx. Melatonin also stimulates the activity of GRd, thereby ensuring the replenishment of GSH⁽²⁹⁾.

By virtue of its ability of generate $\text{NO}\cdot$, nitric-oxide synthase (NOS) can be viewed as a pro-oxidative enzyme. The discovery of the inhibition of NOS by melatonin would directly reduce oxidative damage caused by $\text{NO}\cdot$, as well as its secondary toxicity after it combines with O_2^- to form ONOO^- ⁽³⁰⁾. Additionally 5-lipoxygenase activity is reduced by melatonin⁽³¹⁾.

Antioxidative actions of melatonin

Free radicals and reactive oxygen intermediates indiscriminately damage molecules, often in the immediate vicinity of where they are generated, leading to cellular death via either

apoptosis or necrosis (Fig 2). The destruction inflicted by these agents is usually measured in terms of damaged macromolecules including nuclear and mitochondrial DNA, polyunsaturated fatty acids (PUFA) and proteins.

the ability of melatonin to reduce the oxidation of macromolecules in the nucleus is consistent with the seemingly high concentrations of the indoleamine in nucleoplasm^[34].





Fig 1. Major enzymes involved in removal of hydrogen peroxide (H₂O₂) from cells include catalase and glutathione peroxidase (GPx), both of which metabolize H₂O₂ and hydroperoxides (ROOH) to nontoxic products. In metabolizing these molecules, glutathione (GSH) is converted to its disulfide form (GSSG) which is enzymatically metabolized back to GSH by the enzyme glutathione reductase (GRd) (not shown). Both GPx and GRd are stimulated by melatonin. If H₂O₂ is not removed enzymatically, it can be converted, via the Fenton reaction, to the highly toxic hydroxyl radical (·OH).

Fig 2. Molecular debris that accumulates within cells as a result of persistent bludgeoning of macromolecules by free radicals and reactive oxygen intermediates eventually leads to cell death due to necrotic processes or apoptosis. While the most toxic agent is generally considered to be the hydroxyl radical (·OH), other agents also participate in this destructive onslaught.

Either endogenously-derived or exogenously-administered melatonin affords nuclear DNA considerable protection from oxygen toxicity^[32]. In 1993, shortly after melatonin's ability to neutralize the ·OH was uncovered, the indole was found to protect hepatic DNA from damage induced by the administration of the chemical carcinogen safrole^[33]. While in this study exogenously administered melatonin was provided, the same group also found that pinealectomized rats (which are deficient in endogenously produced melatonin) accumulate oxidatively damaged DNA more rapidly than intact rats when given safrole, thus illustrating that melatonin is not only pharmacologically relevant as an anti-oxidant but physiologically as well. Certainly,

Since these early observations, the ability of melatonin to abate nuclear DNA damage normally induced by free radicals has been repeatedly confirmed. Ionizing radiation, a physical agent known to cause the homolytic scission of H₂O to generate the ·OH, is highly carcinogenic because of its destructive actions at the level of the genome. Since free radicals are involved in ionizing radiation-induced damage, we used this model to test the protective actions of melatonin. Using a variety of cytogenetic procedures to estimate chromosomal damage, giving melatonin in advance of exposure to ionizing radiation protects the cells from genetic damage; this was shown in both *in vivo* and *in vitro* studies and in human and nonhuman cells^[35,36].

In the last 3 years numerous experiments have been reported illustrating that melatonin readily enters the nucleus, situates itself near DNA and shelters the genetic material from oxidative attack^[37]. Considering that, for example, the ·OH, once generated, is estimated to travel no more than 0.3 – 0.5 nm before it interacts with another molecule such as DNA, the melatonin in the nucleus must be in very close proximity of the DNA, ie, within the reacting distance of the ·OH.

By far the largest number of publications related to the protective effects of melatonin against free radicals have measured products of lipid peroxidation^[38]. There are several reasons for this. Firstly, lipids are easily oxidized so their damaged products often appear in relatively high concentrations. Secondly, there are a number of simple and reliable assays to measure the products of lipid peroxidation. And thirdly, the mechanisms of oxidative damage to lipids are generally well understood. While melatonin has clearly been shown capable of reducing oxidative damage to membrane lipids in a large number of experimental situations^[39-41], there seems to be some disagreement as to how this beneficial effect is achieved. Whereas early claims indicated that melatonin was a powerful chain breaking antioxidant (like vitamin E)^[13], later studies have not always come to the same conclusion^[14,15]. Thus, rather than preventing excessive lipid damage by interrupting the chain reaction of events that occur during the breakdown of lipids by scavenging the ROO \cdot , the primary means by which melatonin counteract oxidative attack to PUFA may be by reducing the initiating events in the process; this would be accomplished when melatonin scavenges reactive species such as the \cdot OH and ONOO $^-$. It may be that melatonin inhibits both the initiation as well as the propagation of lipid peroxidation and that the relative importance of each of these processes in protecting PUFA from the destructive effects of reactive species varies under different circumstances.

Because of melatonin's ability to pass through all membranes and apparently be sequestered in all cells and organs, its protective effects against lipid peroxidation are widespread. This contrasts with some other lipid antioxidants, eg, vitamin E, which because of their inability to efficiently penetrate the blood-brain-barrier, are not especially effective in reducing lipid breakdown in the central nervous system (CNS). By contrast, in a variety of experimental settings melatonin, when administered acutely, has been shown to retard damage to the CNS induced by free radical generating agents or toxins^[42-45].

Methods to measure oxidative damage to proteins are less numerous and more difficult to reliably perform compared to the techniques for

measuring lipid and DNA damage. Because this, there are generally fewer reports related to the damaging effects of reactive species on protein. Thus, it is not surprising that melatonin has been less frequently tested for its ability to protect against free radical-induced protein damage. It has been inferred from a number of studies, however, that melatonin reduces protein damage caused by free radicals^[46]. Furthermore, in cell-free *in vitro* studies we have recently shown melatonin to protect purified proteins from the damaging effect of the \cdot OH (unpublished observations by S. Kim, J Cabrera, R J Reiter).

Final Commentary

The unexpected discovery that the chief secretory product of the pineal body, melatonin, functions as a free radical scavenger has led to a vast number of studies in recent years which have clarified some of the multiple actions by which melatonin protects macromolecules from oxidative damage. While melatonin has been shown to have both direct and indirect means for neutralizing free radicals and ROI, there may be other processes by which melatonin assists cells in resisting oxidative damage. As an example, melatonin chelates metals^[47] and acts at the level of the cell membrane to maintain its optimal fluidity^[48,49]. This latter action would ensure the proper functioning of the numerous critical activities of cellular membranes thereby helping them to resist toxic challenges. It is likely that other actions of melatonin will be discovered that permit it to function as an efficient antioxidant.

The multiple actions of melatonin would help to explain how a molecule, which is normally in such low concentrations in the cell (perhaps in the low to medium nanomolar range when it is at its highest levels) is so effective as an antioxidant. The fact that melatonin is normally replenished (it is synthesized perhaps in all aerobic organisms), is also an important feature of its ability to provide protection against oxidants. While melatonin is endogenously produced in young mammals, increasing age is associated with a marked reduction in the generation of this important antioxidant^[50,51].

Since oxidative damage accumulates more rapidly in older individuals, it has been suggested that the

age-associated drop in melatonin may be causative in the acceleration of oxidatively damaged products^[52,53]. This has clear implications for a variety of age-related debilitating conditions, eg, cardiovascular^[54] and neurodegenerative diseases^[55-57] as well as for aging itself^[58] since each of these processes involves free radical damage^[59]. Of particular interest in this regard is Alzheimer disease and associated dementias. Alzheimer disease is believed to be related in part to oxidative destruction of critical neurons^[56]. Recent studies have shown that melatonin may be beneficial in preventing the loss of these cells^[42] and delaying the progression of the disease^[42]. This contention is supported by the recent observation that pinealectomy (which creates a relative melatonin deficiency throughout life) in young rats leads to an accelerated accumulation of oxidatively damaged lipid and DNA especially in the brain (unpublished observations by RJ Reiter, DX Tan, W Qi, SJ Kim). In view of the findings summarized herein, investigations into melatonin's antioxidative actions will likely continue unabated in the foreseeable future.

REFERENCES

- 1 Fridovich I. The biology of oxygen radicals. *Science* 1978; 201: 875-880.
- 2 Pryor WA, Squadrito GL. The chemistry of peroxynitrite, a product of the reaction of nitric oxide with superoxide. *Am J Physiol* 1995; 268: L699-L722.
- 3 Reiter RJ. Pineal melatonin: cell biology of its synthesis and of its physiological interactions. *Endocr Rev* 1991; 12: 151-180.
- 4 Reiter RJ, Melchiorri D, Sewerynek E, Poeggeler B, Barlow-Walden L, Chuang SH, *et al.* A review of the evidence supporting melatonin's role as an antioxidant. *J Pineal Res* 1995; 18: 1-11.
- 5 Costa EJ, Lopes RH, Lamy-Freund MT. Permeability of pure lipid bilayers to melatonin. *J Pineal Res* 1995; 18: 123-6.
- 6 Shida CS, Castrucci AML, Lamy-Freund MT. High melatonin solubility in aqueous medium. *J Pineal Res* 1994; 16: 198-201.
- 7 Ianas O, Olenescu R, Badescu I. Melatonin involvement in oxidative stress. *Rom J Endocrinol* 1991; 29: 147-53.
- 8 Tan DX, Chen LD, Poeggeler B, Manchester LC, Reiter RJ. Melatonin: a potent, endogenous hydroxyl radical scavenger. *Endocrine J* 1993; 1: 57-60.
- 9 Matuszek Z, Reszka KJ, Chignell CF. Reaction of melatonin and related indoles with hydroxyl radical: EPR and spin trapping investigations. *Free Radical Biol Med* 1997; 23: 367-72.
- 10 Susa N, Ueno S, Furukawa Y, Ueda J, Sugiyama M. Potent protective effect of melatonin on chromium (VI)-induced DNA single strand breaks, cytotoxicity, and lipid peroxidation in primary cultures of rat hepatocytes. *Toxicol Appl Pharmacol* 1997; 144: 377-84.
- 11 Stasica P, Ulanski P, Rosiak JM. Melatonin as a hydroxyl radical scavenger. *J Pineal Res* 1998; 25: 65-6.
- 12 Li XJ, Zhang LM, Gu J, Zhang AZ, Sun FY. Melatonin decreases production of hydroxyl radical during cerebral ischemia-reperfusion. *Acta Pharmacol Sin* 1997; 18: 394-96.
- 13 Pieri C, Marra M, Moroni F, Recchioni R, Marcheselli F. Melatonin: a peroxy radical scavenger more effective than vitamin E. *Life Sci* 1994; 55: 271-6.
- 14 Marshall KA, Reiter RJ, Poeggeler B, Aruoma OI, Halliwell B. Evaluation of the antioxidant activity of melatonin *in vitro*. *Free Radical Biol Med* 1996; 21: 307-15.
- 15 Longoni B, Salgo MG, Pryor WA, Marchiafava PL. Effects of melatonin on lipid peroxidation induced by oxygen radicals. *Life Sci* 1998; 62: 853-9.
- 16 Reiter RJ. Antioxidant actions of melatonin. *Adv Pharmacol* 1997; 38: 103-17.
- 17 Cagnoli CM, Atabay C, Kharlamova E, Manev H. Melatonin protects neurons from singlet oxygen-induced apoptosis. *J Pineal Res* 1995; 18: 222-6.
- 18 Gilad E, Cuzzocrea S, Zingarelli B, Salzman AL, Szabo C. Melatonin is a scavenger of peroxynitrite anion. *Life Sci* 1997; 60: PL-169-74.
- 19 Cuzzocrea S, Zingarelli B, Costantino G, Caputi AP. Protective effect of melatonin in a non-septic shock model induced by zymosan in the rat. *J Pineal Res* 1998; 25: 24-33.
- 20 Noda Y, Mori A, Liburdy R, Packer L. Melatonin and its precursors exhibit nitric oxide scavenging activity. *Pathophysiology* 1998; 5 (Suppl 1): 85.
- 21 Guerrero JM, Reiter RJ, Ortiz GG, Pablos MI, Sewerynek E, Chuang JI. Melatonin prevents increases in neural nitric oxide and cyclic GMP production after transient brain ischemia and reperfusion in the Mongolian gerbil (*Meriones unguiculatus*). *J Pineal Res* 1997; 23: 24-31.
- 22 Morrey KM, McLachlan JA, Serkin CD, Bakouche O. Activation of human monocytes by the pineal hormone melatonin. *J Immunol* 1994; 153: 2671-80.
- 23 Pieri C, Recchioni R, Moroni F, Marcheselli F, Marra M, Marinoni S, *et al.* Melatonin regulates the respiratory burst of human neutrophils and their depolarization. *J Pineal Res* 1998; 24: 43-9.
- 24 Hardeland R, Reiter RJ, Poeggeler B, Tan DX. The significance of the metabolism of the neurohormone melatonin: antioxidative protection and formation of bioactive substances. *Neurosci Biobehav Rev* 1993; 17: 347-57.
- 25 Roberts JE, Hu DN, Wishart JF. Pulse radiolysis studies of melatonin and chloromelatonin. *J Photochem Photobiol* 1998; 42: 125-32.
- 26 Kolter M, Rodriguez C, Sainz RM, Antolin I, Menendez-Pelaez A. Melatonin increases gene expression for antioxidant enzymes in rat brain cortex. *J Pineal Res* 1998; 24: 83-9.
- 27 Pablos MI, Reiter RJ, Ortiz GG, Guerrero JM, Agapito MT, Chuang JI, Sewerynek E. Rhythms of glutathione peroxidase and glutathione reductase in brain of chick and their inhibition

- by light. *Neurochem Int* 1998; 32: 69-75.
- 28 Sies H, Klotz LO, Sharov VS, Tanler R, Asmann A, Briviba K. Glutathione peroxidases as peroxynitrite reductases. *Pathophysiology* 1998; 5 (Suppl 1): 61.
- 29 Barlow-Walden LR, Reiter RJ, Abe M, Pablos M, Menendez-Pelaez A, Chen KD, *et al.* Melatonin stimulates brain glutathione peroxidase activity. *Neurochem Int* 1995; 26: 497-502.
- 30 Pozo D, Reiter RJ, Calvo JR, Guerrero JM. Inhibition of cerebellar nitric oxide synthase and cyclic GMP production by melatonin via complex formation with calmodulin. *J Cell Biochem* 1997; 65: 430-42.
- 31 Carlberg C, Weisenberg I. The orphan receptor family RZR/ROR, melatonin and 5-lipoxygenase: an unexpected relationship. *J Pineal Res* 1995; 18: 171-8.
- 32 Reiter RJ, Tan DX, Kim SJ, Qi W. Melatonin as a pharmacological agent against oxidative damage to lipids and DNA. *Proc West Pharmacol Soc*, in press.
- 33 Tan DX, Reiter RJ, Chen LD, Poeggeler B, Manchester LC, Barlow-Walden LR. Both physiological and pharmacological levels of melatonin reduce DNA adduct formation induced by the carcinogen safrole. *Carcinogenesis* 1994; 15: 215-8.
- 34 Finocchiaro LM, Glikin GC. Intracellular melatonin distribution in cultured cell lines. *J Pineal Res* 1998; 24: 22-34.
- 35 Vijayalaxmi, Reiter RJ, Meltz M. Melatonin protects human blood lymphocytes from radiation-induced chromosome damage. *Mutat Res* 1995; 346: 23-31.
- 36 Vijayalaxmi, Reiter RJ, Herman TS, Meltz ML. Melatonin reduces gamma radiation-induced primary DNA damage in human blood lymphocytes. *Mutat Res* 1998; 397: 203-8.
- 37 Reiter RJ. Functional pleiotropy of the neurohormone melatonin: antioxidant protection and neuroendocrine regulation. *Front Neuroendocrinol* 1995; 16: 383-415.
- 38 Reiter RJ, Carneiro RC, Oh CS. Melatonin in relation to cellular antioxidative defense mechanisms. *Horm Metab Res* 1997; 29: 363-72.
- 39 Li ZR, Reiter RJ, Fujimori O, Oh CS, Duan YP. Cataractogenesis and lipid peroxidation in newborn rats treated with buthionine sulfoximine: preventive actions of melatonin. *J Pineal Res* 1997; 22: 117-23.
- 40 Kelly MR, Loo G. Melatonin inhibits oxidative modification of human low-density lipoprotein. *J Pineal Res* 1997; 22: 203-9.
- 41 Princ FG, Juknat AA, Maxit AG, Cardalda C, Battle A. Melatonin's antioxidant protection against δ -aminolevulinic acid-induced oxidative damage in rat cerebellum. *J Pineal Res* 1997; 23: 40-6.
- 42 Pappolla MA, Sos M, Omar RA, Bick RJ, Hickson-Bick DLM, Reiter RJ, *et al.* Melatonin prevents death of neuroblastoma cells exposed to the Alzheimer amyloid peptide. *J Neurosci* 1997; 17: 1683-90.
- 43 Mesenge C, Margail I, Verrecchia C, Allix M, Boulu RG, Plokine M. Protective effect of melatonin in a model of traumatic brain injury in mice. *J Pineal Res* 1998; 25: 41-6.
- 44 Lezoualc'h F, Sparapani M, Behl C. N-acetyl-serotonin (normelatonin) and melatonin protect neurons against oxidative challenges and suppress the activity of the transcription factor NF-kappa B. *J Pineal Res* 1998; 24: 168-78.
- 45 Reiter RJ. Oxidative damage in the central nervous system: protection by melatonin. *Prog Neurobiol* 1998; 56: 359-69.
- 46 Abe M, Reiter RJ, Orhii PB, Hara M, Poeggeler B. Inhibitory effect of melatonin on cataract formation in newborn rats: evidence for an antioxidative role for melatonin. *J Pineal Res* 1994; 17: 94-100.
- 47 Limson J, Nyokong T, Daya S. The interaction of melatonin and its precursors with aluminum cadmium, copper, iron, lead and zinc: an adsorptive voltammetric study. *J Pineal Res* 1998; 24: 15-21.
- 48 Garcia JJ, Reiter RJ, Guerrero JM, Escames G, Yu BP, Oh CS, *et al.* Melatonin prevents changes in microsomal membrane fluidity during induced lipid peroxidation. *FEBS Lett* 1997; 408: 297-300.
- 49 Garcia JJ, Reiter RJ, Ortiz GG, Oh CS, Tang L, Yu BP, Escames G. Melatonin enhances tamoxifen's ability to prevent the reduction in microsomal membrane fluidity induced by lipid peroxidation. *J Membr Biol* 1998; 162: 59-65.
- 50 Reiter RJ, Richardson BA, Johnson LY, Ferguson BN, Dinh DT. Pineal melatonin rhythm: reduction in aging Syrian hamsters. *Science* 1980; 210: 1372-3.
- 51 Reiter RJ, Tan DX, Poeggeler B, Menendez-Pelaez B, Chen LD, Saarela S. Melatonin as a free radical scavenger: implications for aging and age-related diseases. *Ann NY Acad Sci* 1994; 719: 1-12.
- 52 Reiter RJ. Oxygen radical detoxification processes during aging: the functional importance of melatonin. *Aging (Milana)* 1995; 7: 340-51.
- 53 Reiter RJ. Aging and oxygen toxicity: relation to changes in melatonin. *Age* 1997; 20: 201-13.
- 54 Tan DX, Manchester LC, Reiter RJ, Qi W, Kim SJ, El-Sokkary G. (1998) Ischemia/reperfusion induced arrhythmias in the isolated rat heart: prevention by melatonin. *J Pineal Res* 1998; 25: 184-91.
- 55 Reiter RJ, Pie J, Garcia JJ. Oxidative toxicity in models of neurodegeneration: responses to melatonin. *Restr Neurol Neurosci* 1998; 12: 135-42.
- 56 Reiter RJ. Oxidative processes and antioxidative defense mechanisms in the aging brain. *FASEB J* 1995; 9: 526-33.
- 57 Uz T, Giusti P, Franceschini D, Kharlamov A, Manev H. Protective effect of melatonin against hippocampal DNA damage induced by intraperitoneal administration of kainate to rats. *Neuroscience* 1996; 73: 631-6.
- 58 Harman D. Free radical theory of aging. *Mutat Res* 1992; 275: 257-66.
- 59 Reiter RJ, Tang L, Garcia JJ, Muñoz-Hoyos A. Pharmacological actions of melatonin in oxygen radical pathophysiology. *Life Sci* 1997; 60: 2255-71.
- 60 Brusco LI, Marquez M, Cardinali DP. Monozygotic twins with Alzheimer's disease treated with melatonin: a case report. *J Pineal Res*, in press.

褪黑激素抑制氧毒性

Russel J REITER¹, 谭敦究, 齐文波

关键词 褪黑激素; 自由基; 抗氧化剂; 氧化应激; 衰老