

## Inhibitory effects of estradiol on inward rectifier and delayed rectifier $K^+$ currents in guinea pig ventricular myocytes<sup>1</sup>

ZHANG Yi<sup>2</sup>, SONG Li-Lin<sup>3</sup>, GU Shuang-Zhen<sup>3</sup>, LU Shen-Gui<sup>3</sup>, ZHOU Zhao-Nian

(Shanghai Institute of Physiology, Chinese Academy of Sciences, Shanghai 200031, <sup>3</sup>Department of Physiology, Hebei Medical University, Shijiazhuang 050017, China)

**KEY WORDS** estradiol; patch-clamp techniques; myocardium; potassium channels; action potentials

### ABSTRACT

**AIM:** To study the effects of estradiol (Est) on inward rectifier  $K^+$  ( $I_{K1}$ ) and delayed rectifier  $K^+$  ( $I_K$ ) channels in isolated guinea pig ventricular myocytes.

**METHODS:** Using whole cell patch-clamp recording techniques. **RESULTS:** Est  $10 \mu\text{mol} \cdot \text{L}^{-1}$  and  $100 \mu\text{mol} \cdot \text{L}^{-1}$  decreased the action potential duration,  $\text{APD}_{50}$ , from  $(474 \pm 71)$  ms to  $(330 \pm 75)$  ms and  $(229 \pm 67)$  ms ( $n = 7$  cells of 7 guinea pigs,  $P < 0.05$ ), respectively. Est  $100 \mu\text{mol} \cdot \text{L}^{-1}$  also decreased  $\text{APD}_{50}$  from  $(587 \pm 60)$  ms to  $(418 \pm 79)$  ms ( $n = 7$ ,  $P < 0.05$ ). Est inhibited  $I_K$  tail current ( $I_K \cdot \text{tail}$ ) concentration-dependently.  $I_K \cdot \text{tail}$  was depressed 53% ( $n = 5$ ,  $P < 0.05$ ) at  $10 \mu\text{mol} \cdot \text{L}^{-1}$  and 80% ( $n = 5$ ,  $P < 0.01$ ) at  $100 \mu\text{mol} \cdot \text{L}^{-1}$  compared with control. Est  $\geq 10 \mu\text{mol} \cdot \text{L}^{-1}$  blocked  $I_{K1}$ . The maximal inhibition of inward current of  $I_{K1}$  occurred at  $-100$  mV test potential was 49% ( $n = 5$ ,  $P < 0.01$ ) and outward current of  $I_{K1}$  at  $-40$  mV was 72% ( $n = 5$ ,  $P < 0.01$ ). The reverse potential shifted negatively, from  $-70$  to  $-76$  mV. **CONCLUSION:** Est possessed blocking effects on both  $I_{K1}$  and  $I_K$  channels in guinea pig ventricular myocytes.

### INTRODUCTION

Estrogens played a key role in sex difference of

cardiovascular diseases<sup>[1]</sup>. It protected the heart against coronary disease and ischemia in some conditions and acted as a coronary risk factor in another condition<sup>[2-4]</sup>. Est inhibited contraction of isolated rabbit heart, guinea pig papillary muscle and guinea pig ventricular myocytes, antagonized experimental arrhythmia and influenced action potentials (AP) of isolated guinea pig papillary muscle<sup>[5-7]</sup>. The present study was to observe the effects of Est on AP, inward rectifier  $K^+$  current ( $I_{K1}$ ) and delayed rectifier  $K^+$  current ( $I_K$ ) in cardiomyocytes.

### MATERIALS AND METHODS

**Preparation of ventricular myocytes** Single ventricular myocytes from guinea pigs ( $\text{♂}$ ,  $n = 11$ , weighing  $261 \text{ g} \pm 47 \text{ g}$ ) were prepared by enzymatic dissociation<sup>[8]</sup>. Briefly, the heart was rinsed in an oxygenated  $\text{Ca}^{2+}$ -free Tyrode's solution. The aorta was cannulated and the heart was retrogradely perfused on a Langendorff apparatus at  $37^\circ\text{C}$ . A perfusion with  $\text{Ca}^{2+}$ -free Tyrode's solution for 5 min was followed by low  $\text{Ca}^{2+}$  ( $50 \mu\text{mol} \cdot \text{L}^{-1}$ ) Tyrode's solution containing 0.03% collagenase and 1% bovine serum albumin (BSA) for 5 min. The ventricles were cut, minced, and gently triturated with a pipette in the low- $\text{Ca}^{2+}$  Tyrode's solution containing BSA at  $37^\circ\text{C}$  for 10 min. The cells were filtered through 200- $\mu\text{m}$  nylon mesh, resuspended in the Tyrode's solution in which the  $\text{Ca}^{2+}$  concentration gradually increased to  $1.0 \mu\text{mol} \cdot \text{L}^{-1}$ . Only the cells with rod shaped and clear cross striation were used for experiments.

**Chemicals and solutions** Est was purchased from Sigma Co and dissolved in ethanol to make a  $10 \text{ mmol} \cdot \text{L}^{-1}$  stock solution. The maximal concentration of ethanol in perfusate was 0.2%. All of BSA, collagenase type II, taurine, HEPES, egtazic acid,

<sup>1</sup> Project supported by the Hebei Science Foundation, No 395356.

<sup>2</sup> Correspondence to Dr ZHANG Yi.

Phn 86-21-6431-3251, ext 8568. Fax 86-21-6474-6305.

E-mail henryzha@sunm.shnc.ac.cn

Received 1998-06-16

Accepted 1998-12-10

Na<sub>2</sub>ATP and K<sub>2</sub>ATP were products of Sigma. 3-(*N*-morpholino)-propanesulfonic acid (MOPS) was purchased from Shanghai Boao Biotech Co. The composition of the Ca<sup>2+</sup>-free Tyrode's solution was: NaCl 100, KCl 10, NaH<sub>2</sub>PO<sub>4</sub> 1.2, MgSO<sub>4</sub> 5.0, glucose 20, taurine 10, MOPS 10 mmol·L<sup>-1</sup>; pH was adjusted with KOH to 7.2. Test solution for AP recording was composed of: NaCl 137, KCl 5.4, MgCl<sub>2</sub> 1.0, CaCl<sub>2</sub> 1.8, HEPES 10, glucose 20 mmol·L<sup>-1</sup>; pH was adjusted with KOH to 7.4. Test solution for K<sup>+</sup> currents recording: AChCl 137, KCl 5.4, MgCl<sub>2</sub> 1.0, HEPES 10, glucose 10 mmol·L<sup>-1</sup>; pH was adjusted with KOH to 7.4. The electrode internal solution for AP recording: KCl 140, MgCl<sub>2</sub> 2.0, egtazic acid 2.0, HEPES 5.0, Na<sub>2</sub>ATP 4.0 mmol·L<sup>-1</sup>; pH was adjusted with KOH to 7.4. The electrode internal solution for K<sup>+</sup> currents recording: KCl 140, MgCl<sub>2</sub> 0.5, egtazic acid 10, HEPES 10, K<sub>2</sub>ATP 5.0 mmol·L<sup>-1</sup>; pH was adjusted with KOH to 7.4.

**Potential and currents recording** Myocytes were placed in a 500 μL chamber on stage of inverted microscope (Olympus CK2). The chamber was continuously superfused with test solutions 2 mL·min<sup>-1</sup> at 25 °C. Membrane currents and AP were recorded using the whole-cell patch-clamp techniques with a patch-clamp amplifier (CEZ 2300, Nihon Kohden, Japan)<sup>[9]</sup>. Patch electrodes were pulled with a vertical puller (PB-7, Narishige, Tokyo, Japan) and had a resistance of 2 - 3 MΩ when filled with electrode internal solution. After gigaseal produced and patch ruptured, AP was recorded in current clamp mode and currents were recorded in voltage clamp mode. Experimental protocols, data acquisition and storage

were accomplished with Pclamp 5.6 (Axon Instrument, USA) running on a personal computer.

**Statistics** Data were expressed as  $\bar{x} \pm s$  and compared with the paired *t*-test.

## RESULTS

**AP** AP in guinea pig ventricular myocytes was evoked by a step current pulse of 90 pA, 10 ms duration at the frequency of 1 Hz. Est 10 μmol·L<sup>-1</sup> decreased APD<sub>50</sub>, from (474 ± 71) ms to (330 ± 75) ms (*n* = 7 cells of seven guinea pigs, *P* < 0.05). Est 100 μmol·L<sup>-1</sup> decreased APD<sub>50</sub> to (229 ± 67) ms (*n* = 7, *P* < 0.01) and APD<sub>90</sub> from (587 ± 60) ms to (418 ± 79) ms (*P* < 0.05). The effects were not canceled completely after Est was washed out. RP and other parameters of AP had no significant change (Tab 1). Ethanol 0.2 % had no significant effect on AP, RP as well as currents.

**I<sub>K</sub> · tail** I<sub>K</sub> · tail in guinea pig ventricular myocytes was obtained by a depolarizing step pulse from the holding potential (E<sub>h</sub>) of -40 mV to 30 mV at the frequency of 0.2 Hz. The step pulse duration was 5 s. Est inhibited I<sub>K</sub> · tail concentration-dependently, 53 % (from 247 ± 51 to 117 ± 27 pA, *n* = 5 cells of 4 guinea pigs, *P* < 0.05) at 10 μmol·L<sup>-1</sup> and 80 % (from 248 ± 51 to 57 ± 13 pA, *P* < 0.01) at 100 μmol·L<sup>-1</sup>. After washout of Est, the inhibition of I<sub>K</sub> · tail was partly recovered (from 57 ± 13 to 84 ± 24 pA) (Fig 1).

**I<sub>K1</sub>** I<sub>K1</sub> was elicited by a number of step pulses (300 ms) from the E<sub>h</sub> of -40 mV to test potential between -100 and +30 mV with step 10 mV. Addition of Est 10 μmol·L<sup>-1</sup> induced a remarkable

**Tab 1.** Effects of Est on AP. *n* = 7 ventricular myocytes of 7 guinea pigs.  $\bar{x} \pm s$ .  
<sup>a</sup>*P* > 0.05, <sup>b</sup>*P* < 0.05, <sup>c</sup>*P* < 0.01 vs control.

	RP/mV	APA/mV	OS/mV	V <sub>max</sub> /V·s <sup>-1</sup>	APD <sub>50</sub> /ms	APD <sub>90</sub> /ms
Solvent	-82 ± 5 <sup>a</sup>	150 ± 18 <sup>a</sup>	68 ± 21 <sup>a</sup>	38 ± 7 <sup>a</sup>	439 ± 56 <sup>a</sup>	558 ± 62 <sup>a</sup>
Control	-81 ± 6 <sup>a</sup>	145 ± 10 <sup>a</sup>	64 ± 14 <sup>a</sup>	33 ± 5 <sup>a</sup>	474 ± 71 <sup>a</sup>	587 ± 60 <sup>a</sup>
Est/μmol·L <sup>-1</sup>						
1	-82 ± 6 <sup>a</sup>	148 ± 17 <sup>a</sup>	65 ± 18 <sup>a</sup>	32 ± 8 <sup>a</sup>	407 ± 32 <sup>a</sup>	522 ± 45 <sup>a</sup>
10	-83 ± 5 <sup>a</sup>	145 ± 18 <sup>a</sup>	62 ± 19 <sup>a</sup>	30 ± 8 <sup>a</sup>	330 ± 75 <sup>a</sup>	460 ± 95 <sup>a</sup>
100	-82 ± 7 <sup>a</sup>	140 ± 28 <sup>a</sup>	58 ± 23 <sup>a</sup>	20 ± 12 <sup>a</sup>	229 ± 67 <sup>c</sup>	418 ± 79 <sup>b</sup>
Washout	-85 ± 2 <sup>a</sup>	149 ± 15 <sup>a</sup>	64 ± 15 <sup>a</sup>	21 ± 6 <sup>a</sup>	294 ± 65 <sup>b</sup>	495 ± 95 <sup>a</sup>




Fig 1. Block of  $I_K$  tail by Est in guinea pig ventricular myocytes. A) Current tracings from a representative cell. B) concentration-dependent block of  $I_K$  tail by Est.  $n = 5$ . <sup>b</sup> $P < 0.05$ , <sup>c</sup> $P < 0.01$  vs control.

depression of  $I_{K1}$ . The inward currents of  $I_{K1}$ , at  $E_i$  -100 mV and -90 mV, were depressed 36 % (from -652 pA  $\pm$  54 pA to -420 pA  $\pm$  60 pA,  $P < 0.01$ ) and 44 % (from -321 pA  $\pm$  53 pA to -178 pA  $\pm$  24 pA,  $P < 0.05$ ) of the control, respectively and outward currents, at  $E_i$  of -40 mV, 29 % (from 126 pA  $\pm$  25 pA to 90 pA  $\pm$  23 pA,  $P < 0.05$ ,  $n = 5$  cells of 5 guinea pigs). Est 100  $\mu\text{mol}\cdot\text{L}^{-1}$  had a more remarkable blocking effect on  $I_{K1}$ . The inward currents of  $I_{K1}$  at  $E_i$  of -100 mV and -90 mV were depressed respectively 49 % (from -652 pA  $\pm$  54 pA to -334 pA  $\pm$  78 pA,  $P < 0.01$ ) and 55 % (from -321 pA  $\pm$  53 pA to -144 pA  $\pm$  27 pA,  $P < 0.05$ ) and outward currents, at  $E_i$  of -60 mV, -50 mV and -40 mV, 43 % (from 150 pA  $\pm$  39 pA to 91 pA  $\pm$  20 pA,  $P < 0.05$ ), 55 % (from 163 pA  $\pm$  23 pA to 72 pA  $\pm$  20 pA,  $P < 0.05$ ) and 72 % (from 126 pA  $\pm$  25 pA to 38 pA  $\pm$  11 pA,  $P < 0.05$ ) of control ( $n = 5$ , Fig 2).

## DISCUSSION

Effects of estrogen on cardiovascular system are




Fig 2. Effects of Est on  $I_{K1}$ .  $n = 5$  ventricular myocytes of 5 guinea pigs. <sup>b</sup> $P < 0.05$ , <sup>c</sup> $P < 0.01$  vs control.

diverse, irregular and even contradictory with each other in some condition<sup>[10]</sup>, which means the mechanism of estrogen is very complex. Although estrogen receptors were found in heart<sup>[11]</sup> and the direct effects of Est on heart were observed, the mechanism of the hormone is not clear yet. In the present study, we firstly demonstrated the inhibitory effects of Est on  $I_{K1}$  and  $I_K$  in single ventricular myocytes of guinea pig in a concentration-dependent manner. It suggested that the inhibitory effects of Est on  $K^+$  channels might play an important role in the effects of Est on heart.

$I_{K1}$  and  $I_K$  channels of myocardium are very important in maintaining normal RP and APD in electrophysiology of heart. Some reported that the blockade of  $K^+$  channels could protect against arrhythmia<sup>[12]</sup>. The inhibitory effects of Est on  $I_{K1}$  and  $I_K$  may be another ionic basis of antagonizing experimental arrhythmia, besides  $I_{Ca}$  channel blocking<sup>[13]</sup>. In the experiment, APD decreasing and  $K^+$  channels blocking seems contradictory because blockade of  $K^+$  channels usually cause the APD shortening. Some reported that Est has a  $\text{Ca}^{2+}$  blocking effect<sup>[13]</sup> and the results of study in our laboratory definitely supported that. It suggested that Est blocked  $\text{Ca}^{2+}$  channel stronger than  $K^+$  channels at the same time, which might be the main cause of APD shortening in myocytes. Interestingly, Est shortened APD in single cardiomyocytes<sup>[13,14]</sup>, but in isolated papillary muscle we previously used<sup>[7]</sup>, Est prolonged the APD significantly. Different effect of Est on APD

between papillary muscles and myocytes has not been understood yet and should be paid attention to.

REFERENCES

1 Morise AP, Dalal JN, Duval RD. Value of simple measure of estrogen status for improving the diagnosis of coronary artery disease in women. *Am J Med* 1993; 94: 491-6.

2 Collins P, Rosano GMC, Jiang C, Lindsay D, Sarrel PM, Poole-Wilson PA. Cardiovascular protection by oestrogen: a calcium antagonist effects? *Lancet* 1993; 341: 1264-5.

3 Barta E, Strec V, Styk J, Okolicany J, Rajecova O. Protective effect of oestradiol on the heart of rats exposed to acute ischaemia. *Physiol Bohemoslov* 1989; 38: 193-200.

4 Myrup B, Jensen GF, McNair P. Cardiovascular risk factors during estrogen-norethindrone and cholecalciferol treatment. *Arch Intern Med* 1992; 152: 2265-8.

5 Raddino R, Poli E, Pela G, Manca C. Action of steroid sex hormones on the isolated rabbit heart. *Pharmacology* 1989; 38: 185-90.

6 Zheng ZZ. The effects of sex hormone on cardiovascular system. *Basic Med Sci Clin* 1987; 7: 16-21.

7 Zhang Y, Hao YC, Song LL, Guo SM, Gu SZ, Lu SG. Effects of sex hormones on action potential and contraction of guinea pig papillary muscle. *Acta Pharmacol Sin* 1998; 19: 248-50.

8 Farmer BB, Mancina M, Williams ES, Watanabe AM. Isolation of calcium tolerance myocytes from adult rat hearts; review of the literature and description of a method. *Life Sci* 1983; 33: 1-18.

9 Hamill OP, Marty A, Neher E, Sakmann B, Sigworth FJ. Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflügers Arch* 1981; 391: 85-100.

10 Zheng ZZ, He LP, Li CQ, Shao ZH, Li YJ, Song SJ, et al. Protection of anerobic heart *in vitro* by estradiol in its physiological level. *Chin J Cardiol* 1989; 17: 45-47.

11 Stumpf WE, Sar M, Aumüller G. The heart; a target organ for estradiol. *Science* 1977; 196: 319-21.

12 Fermim B, Jurkiewicz NK, Jow B, Guinasso PJ Jr, Baskin EP, Lynch JJ Jr, et al. Use-dependent effects of the class III antiarrhythmic agent NE-10064 (Azimilide) on cardiac repolarization; block of delayed rectifier potassium and L-type calcium currents. *J Cardiovasc Pharmacol* 1995; 26: 259-71.

13 Jiang C, Poole-Wilson PA, Sarrel PM, Mochizuki S, Collins P, MacLeod KT. Effects of 17β-estradiol on

contraction, Ca<sup>2+</sup> current and intracellular free Ca<sup>2+</sup> in guinea-pig isolated cardiac myocytes. *Br J Pharmacol* 1992; 106: 739-45.

14 Wang XQ, Jiang Y, Zhong GG, Yang GY. Influence of estradiol benzoate on the electric activity of primary cultured rat heart cells. *Chin J Endemiol* 1990; 9: 347-49.

雌二醇抑制豚鼠心室肌细胞内向整流和延迟整流钾通道电流<sup>1</sup>

R965.2

张翼<sup>2</sup>, 宋立林<sup>3</sup>, 谷双振<sup>3</sup>, 卢慎圭<sup>3</sup>, 周兆年 (中国科学院上海生理研究所, 上海 200031; <sup>3</sup>河北医科大学生理教研室, 石家庄 050017, 中国)

关键词 雌二醇; 膜片箱技术; 心肌; 钾通道; 动作电位

药理

目的: 研究雌二醇(Estradiol, Est)对心室肌细胞动作电位(AP)、内向整流钾通道电流( $I_{K1}$ )及延迟整流钾通道电流( $I_K$ )的影响。方法: 全细胞膜片箱技术。结果: EST 10  $\mu\text{mol}\cdot\text{L}^{-1}$ 使豚鼠心室肌细胞 AP 时程明显缩短, APD<sub>50</sub>由给药前(474 ± 71) ms 缩短至(330 ± 75) ms ( $P < 0.05$ ), Est 100  $\mu\text{mol}\cdot\text{L}^{-1}$ 使 APD<sub>50</sub>缩短至(229 ± 67) ms ( $P < 0.01$ ), 使 APD<sub>90</sub>由(587 ± 60) ms 缩短至(418 ± 79) ms ( $P < 0.05$ )。Est 浓度依赖性地抑制  $I_K$  尾电流( $I_K\cdot\text{tail}$ ), 10  $\mu\text{mol}\cdot\text{L}^{-1}$ 浓度下,  $I_K\cdot\text{tail}$  减少 53 % ( $P < 0.05$ ), 100  $\mu\text{mol}\cdot\text{L}^{-1}$ 浓度下,  $I_K\cdot\text{tail}$  减少 80 % ( $P < 0.05$ )。10  $\mu\text{mol}\cdot\text{L}^{-1}$ 以上浓度 Est 明显抑制  $I_{K1}$ , 在 -100 mV 刺激电压下, 内向电流最大抑制为 49 % ( $P < 0.01$ ); 在 -40 mV 刺激电压下, 外向电流最大抑制为 72 % ( $P < 0.01$ )。同时, Est 使  $I_{K1}$  翻转电位向负电位方向移位(由 -70 mV 变为 -76 mV)。结论: Est 对豚鼠心室肌细胞  $I_{K1}$  和  $I_K$  通道具有明显的抑制作用。

(责任编辑 李颖)