

Blocking L-calcium current by *l*-tetrahydropalmatine in single ventricular myocyte of guinea pigs

HUANG Kai¹, DAI Gui-Zhu, LI Xin-Hua, Fan Qu², CHENG Lan², Feng Yi-Bai, XIA Guo-Jin², YAO Wei-Xin²
(Department of Cardiology, Cardiovascular Disease Institute, Union Hospital;

²Department of Pharmacology, Basic Medical College, Tongji Medical University, Wuhan, 430022, China)

KEY WORDS tetrahydropalmatine; patch-clamp techniques; calcium current; myocardium; cultured cells; membrane potentials

ABSTRACT

AIM: To study the effect of *l*-tetrahydropalmatine (*l*-THP) on L-type calcium channel. **METHODS:** Patch clamp technique (whole cell recording) was used to record L-Ca²⁺ current in single cardiac myocyte. **RESULTS:** 1) *l*-THP 1, 10, and 100 μmol · L⁻¹ reduced *I*_{Ca-max} from (999 ± 93) pA to (700 ± 111) pA, (582 ± 66) pA, and (420 ± 112) pA (*n* = 6, *P* < 0.01), respectively. 2) *l*-THP reduced the voltage at half-maximal inactivation (*V*_{1/2}) of L-Ca²⁺ channel to more negative potentials by 9 mV (*n* = 5, *P* < 0.05). 3) *l*-THP caused both tonic and use-dependent reduction of Ca²⁺ current. Tonic block of *l*-THP on Ca²⁺ current was 46 % ± 8 % (*n* = 6, *P* < 0.01). The degree of use dependent blocking was 13.5 % ± 2.4 % (*n* = 6, *P* < 0.05) at 1 Hz, the degree increased to 44 % ± 5 % (*n* = 6, *P* < 0.01) at 3 Hz. 4) *l*-THP delayed half-recovery time of Ca²⁺ channel recovery from inactivity from (94 ± 39) ms to (170 ± 42) ms (*n* = 6, *P* < 0.01). **CONCLUSION:** *l*-THP has a moderate inhibitory effect on L-Ca²⁺ current.

INTRODUCTION

l-Tetrahydropalmatine (*l*-THP) is an alkaloid extracted from *Corydalis ambigua* Cham. Previous

researches had documented its abilities of anti-arrhythmia and L-Ca²⁺ current blocking effects^[1,2]. However, the characteristics of its effect on L-calcium current is still unknown. Therefore, the present experiments were carried out to reveal them in single guinea pig ventricular myocyte.

MATERIALS AND METHODS

Drugs *l*-THP was made by the Nanning Second Pharmaceutical Factory. *l*-THP stocking solution (10 mmol · L⁻¹) was prepared with distilled water and the pH was adjusted to 7.0 with HCl before use. Tyrode's solution contained: NaCl 137, CsCl 5.4, CaCl₂ 1.8, MgCl₂ 1.0, NaHCO₃ 1.0, NaH₂PO₄ 0.05, glucose 5.55, HEPES 10 mmol · L⁻¹, the pH was adjusted to 7.35 with NaOH. Calcium free Tyrode's solution was Tyrode's solution without CaCl₂. The ventricular myocyte dissociation solution was calcium free Tyrode's solution containing collagenase 0.33 g · L⁻¹ (Sigma), BSA 0.25 g · L⁻¹ (Sigma), protease E 0.25 g · L⁻¹ (Sigma). Intracellular solution consisted of: CsCl 120, NaCl 10, CaCl₂ 1.0, Mg-ATP 5.0, egtazic acid 11 (Sigma), HEPES 10 (Sigma) mmol · L⁻¹, pH was adjusted to 7.3 with CsOH.

Preparation of single ventricular myocyte

Male or female guinea pigs weighing (300 ± 40) g (species: England undercoat guinea pigs; Grade II; Certificate number: 19-023; Medical Experimental Animal Center of Tongji Medical University) were used. Single ventricular cells of guinea pig were obtained by an enzymatic dissociation method^[3]. First, the heart was perfused with dissociation solution through coronary arteries for 5 min at 37.5 °C. Subsequently, the dissociation solution was washed out by calcium free Tyrode's solution. Then several pieces of ventricular tissue were dissected and gently agitated

¹ Correspondence to Dr HUANG Kai.

Phn 86-27-8572-6696. Fax 86-27-8380-8080.

E-mail cjhuang@public.wuhan.cngb.com

Received 1998-09-28

Accepted 1999-04-12

in Tyrode's solution contained Ca^{2+} $0.2 \text{ mmol}\cdot\text{L}^{-1}$. After that the isolated ventricular myocytes were stored in Tyrode's solution for 2 h before use.

Electric recordings The isolated cells were set to a 1.0 mL chamber mounted on the stage of an inverted microscope. After being settled to the bottom of the chamber, cells were superfused with Tyrode's solution to wash out the dead cells. The experiments were performed at room temperature (21 ± 2) $^{\circ}\text{C}$.

Micropipettes whose tips were 1–2 μm were constructed from stardore capillary tubes and were heat-polished to have resistances of 1–3.5 $\text{M}\Omega$ when they were filled with the internal solution and immersed in the Tyrode's solution. Recordings were made in the whole cell voltage-clamp configuration of the patch-clamp technique with a PC-2 patch clamp amplifier. DA-1 data collection and analysis system was used to record and analyse I_{Ca} . Sampled data were stored on the hard disk of computer for later analysis. To investigate the effect of *l*-THP on the voltage dependent of Ca^{2+} current availability, the twin-pulse inactivation protocol were used to evaluate effects of membrane potential on the peak Ca^{2+} current of a subsequent depolarization (*f*-curve) with and without *l*-THP. The holding potential was maintained at -40 mV , and conditioning pulses (4 s) to potentials between -60 and $+20 \text{ mV}$ were applied (Fig 2, inset)^(4,5). Peak Ca^{2+} current was measured during a subsequent test pulse and plotted as a function of the prepulse membrane potential. Prepulses to more positive potentials result in inactivation of Ca^{2+} channel^(6,7). The data presented were recored after a 10-min drug exposure. The frequency of stimulation was 0.10 Hz.

In order to observe the use-dependent effect of *l*-THP on Ca^{2+} current, different frequency (1 Hz, 3 Hz) stimulus was used to activate Ca^{2+} current⁽⁸⁾. The membrane potential was held at -40 mV without pulsing for 5 min before each train. The command potential was $+20 \text{ mV}$ followed by a depolarization to -40 mV with 10 ms. Tonic block was calculated as a diminution in the peak I_{Ca} of the first depolarizing pulse after exposure to *l*-THP compared with control⁽⁹⁾. Twin-pulse was used to study *l*-THP on recovery time from inactivity of Ca^{2+} channel. Holding potential maintained -80 mV , prepulse and test-pulse both were 10 mV, interval of two pulse was from 5 to 1280 ms (5, 10, 20, 40, 80, 1280).

Statistic analysis Values were expressed as $\bar{x} \pm s$ by *t* test.

RESULTS

Ca^{2+} current When the holding potential was -40 mV and the cells were voltage-clamped from -40 mV to different membrane potential (-30 to $+60 \text{ mV}$) at a frequency of 0.10 Hz, the inward I_{Ca} was invoked. When the command potential was 10 mV, it reached its maximum. After adding *l*-THP 1, 10, and 100 $\mu\text{mol}\cdot\text{L}^{-1}$, respectively, $I_{\text{Ca-max}}$ reduced from (999 ± 93) pA to (700 ± 111) pA, (582 ± 66) pA, and (420 ± 112) pA ($n = 6, P < 0.01$), respectively (Fig 1A).




Fig 1. A) *l*-THP on Ca^{2+} current of guinea pig single ventricular cell. B) Example of Ca^{2+} current with (left) or without (right) *l*-THP $1 \mu\text{mol}\cdot\text{L}^{-1}$ in Tyrode's solution. C) *I*–*V* relationship curves of L-Ca^{2+} . ■; Control. □; *l*-THP $1 \mu\text{mol}\cdot\text{L}^{-1}$. ●; 10 $\mu\text{mol}\cdot\text{L}^{-1}$. ○; 100 $\mu\text{mol}\cdot\text{L}^{-1}$.




Fig 2. Effect of *l*-THP ($10 \mu\text{mol} \cdot \text{L}^{-1}$) on voltage-dependence of calcium current. $n = 5$ Cells from 5 guinea pigs. $\bar{x} \pm s$.

After *l*-THP was washed out, the Ca^{2+} current reversed (Fig 1B). The data obtained from the experiments fitted with the Boltzmann equation $f = 1 / (1 + \text{EXP} [(V_{\text{pre}} - V_{1/2})/k])$. The free parameters, $V_{1/2}$ and k , quantify the half-maximal potential where current was half-maximal and the slope of activation curves ($d\infty$). The $V_{1/2}$ values in control and in *l*-THP $10 \mu\text{mol} \cdot \text{L}^{-1}$ were 7.12 mV and 8.33 mV ($P > 0.05$), respectively. The k values in control and in *l*-THP $10 \mu\text{mol} \cdot \text{L}^{-1}$ were -7.11 mV and -6.90 mV ($P > 0.05$), respectively.

Voltage dependent of Ca^{2+} current availability Under control condition, the amplitude of Ca^{2+} current was the largest after a prepulse to potential more negative than -50 mV . *l*-THP decreased Ca^{2+} current at -60 to $+20 \text{ mV}$, and the voltage at half-maximal inactivation ($V_{1/2}$) was reduced to more negative potentials by 9 mV [$(-29 \pm 5) \text{ mV}$ vs $(-38 \pm 7) \text{ mV}$, $n = 5$, $P < 0.05$] (Fig 2).

Use dependent *l*-THP caused a marked tonic block of Ca^{2+} channels. Ca^{2+} current was reduced during the first pulse of the train. The tonic block of *l*-THP $10 \mu\text{mol} \cdot \text{L}^{-1}$ on Ca^{2+} current was $46\% \pm 8\%$ ($n = 6$, $P < 0.01$). After exposure to *l*-THP $10 \mu\text{mol} \cdot \text{L}^{-1}$, compared with that in control, the degree of use dependent blocking was $13.5\% \pm 2.4\%$ ($n = 6$, $P < 0.05$) at 1 Hz , the degree increased to $41\% \pm 5\%$ ($n = 6$, $P < 0.01$) at 3 Hz . The blocking effect

of 3 Hz was stronger than that of 1 Hz stimulus ($n = 6$, $P < 0.01$) (Fig 3).




Fig 3. Use-dependent block of Ca^{2+} current by *l*-THP $10 \mu\text{mol} \cdot \text{L}^{-1}$. Ca^{2+} current at the first stimulus in control (A) and in *l*-tetrahydropalmitate (B) at 3 Hz , frequency-dependent inhibition curve of Ca^{2+} current (C). $n = 6$ cells from 6 guinea pigs. $\bar{x} \pm s$.

Recovery from inactivity of Ca^{2+} channel

l-THP delayed Ca^{2+} channel half-recovery time from (94 ± 39) to $(170 \pm 42) \text{ ms}$ ($n = 6$, $P < 0.01$) (Fig 4).

DISCUSSION

l-THP can reduce the L- Ca^{2+} current in the ventricular myocytes of guinea pig. This effect seems stronger than that of berberine on L- Ca^{2+} current^[10]. Another important result in this experiment was that *l*-THP blocked cardiac L-calcium channel potently at which most channels were inactivated. Just as Fig 2 shown, when the 4-s conditioning pulses were applied to different voltages, the resulting steady-state inactivation curve was shift to the hyperpolarizing direction by 9 mV ($n = 5$, $P < 0.01$), this result showed that its effect on the inactivation of Ca^{2+} channel was the same as that of berberine^[10]. The result which is characterised by a number of Ca^{2+} channel ligands




Fig 4. Effect of *l*-THP on recovery time from inactivation of Ca^{2+} channel. $n = 6$ cells from 6 guinea pigs. $\bar{x} \pm s$.

suggested that *l*-THP should have the affinity to the inactivated state of the Ca^{2+} channel. The modulated receptor model could account for the results described in this paper^[11]. The more the channels were bound by *l*-THP, the more the Ca^{2+} current was decreased. This consisted with the concentration-dependent action of *l*-THP on Ca^{2+} channel.

Use dependence and frequency dependence are the common characters of Ca^{2+} channel blockers. *l*-THP had these characters also. Its Ca^{2+} channel blocking effect was significantly stronger at 3 Hz than that at 1 Hz stimulus. But berberine has no effect^[10]. This character is very useful in the therapy of tachycardia when the heart rate is very fast. Another finding in this experiment was that *l*-THP could delay Ca^{2+} channel recovery time from inactivity which means that Ca^{2+} channel would spend much more time to recover. Namely, the duration of inactivation was added. This character should contribute to its blocking effect on Ca^{2+} channel.

We observed a tonic component of block by *l*-THP, which could result from either a specific receptor-mediated inhibition on the channel protein as suggested by the modulated receptor hypothesis for ion channel blockers, or from some indirect or nonspecific action. Our results did not allow us to discriminate between these possibilities.

"Run down" of Ca^{2+} current had ever been reported in whole cell configuration of patch clamp

technique. In this experiment, many methods were used to decrease this phenomenon: 1) ATP and egtazic acid were maintained in the pipette solution. 2) The strict selection of cells. After Ca^{2+} current was evoked, it was observed for 10 - 15 min, if the reduction was less than 10 %, then the drug was added, or the cell was given up otherwise. 3) Washing-out. the Ca^{2+} current reduction after *l*-THP was added could recover after drug was washed out by Tyrode's solution. 4) The lessening of I_{Ca} caused by the run-down in control was less than 10 % throughout the observation period, which has no significant influence on results, so that we could consider that the decrease in I_{Ca} was really caused by *l*-THP, not by the "run down" phenomenon^[12].

In summary, the inhibition of Ca^{2+} current caused by *l*-THP was use-dependent and membrane potential-dependent, meanwhile, the time of calcium channel recovery from inactivation was delayed by *l*-THP. These suggest that *l*-THP has a moderate inhibitory effect on L-Ca^{2+} current. On the other hand, these results can be very helpful in the therapy of cardiac arrhythmia with *l*-THP.

REFERENCES

- 1 Zhao DK, Zhao GS, Qiu PL. The effect of *l*-tetrahydropalmatine on experimental arrhythmia. *Acta Acad Med Xi-an* 1985; 6: 332-3.
- 2 Wang DJ, Mao HY, Lei M, Fang YZ, Tu YS. *l*-Tetrahydropalmatine in the treatment of atrial fibrillation. *Chin Integ Trad West Med* 1993; 13: 455-7.
- 3 Iijima T, Irisawa H, Karneyama M. Membrane currents and their modification by acetylcholine in isolated single atrial cells of the guinea-pig. *J Physiol (Lond)* 1985; 359: 485-501.
- 4 Rampe D, Skattebol A, Triggle DJ, Brown AM. Effects of McN-6186 on voltage-dependent Ca^{2+} channels in heart and pituitary cells. *J Pharmacol Exp Ther* 1989; 248: 164-70.
- 5 Cerbai E, Giotti A, Mugelli A. Characteristics of L-type calcium channel blockade by lacidipine in guinea-pig ventricular myocytes. *Br J Pharmacol* 1997; 120: 667-75.
- 6 Beeler GW Jr, Reuter H. Membrane calcium current in ventricular myocardial fibres. *J Physiol (Lond)* 1970; 207: 191-209.
- 7 Isenberg G, Klockner U. Calcium currents of isolated bovine ventricular myocytes are fast and of large amplitude. *Pflügers Arch* 1982; 395: 30-41.

- 8 Valenzuela C, Bennett PB. Voltage- and use-dependent modulation of calcium channel current in guinea pig ventricular cells by amiodarone and des-oxo-amiodarone. *J Cardiovasc Pharmacol* 1991; 17: 894-902.
- 9 Yamane T, Sunami A, Sawanobori T, Hiraoka M. Use-dependent block of Ca^{2+} current by nifedipine in guinea-pig ventricular myocytes: a possible ionic mechanism of action potential shortening. *Br J Pharmacol* 1993; 108: 812-8.
- 10 Xu SZ, Zhang Y, Ren JY, Zhou ZN. Effects of berberine on L- and T-type calcium channels in guinea pig ventricular myocytes. *Acta Pharmacol Sin* 1997; 18: 515-8.
- 11 Bean BP. Nitrendipine block of cardiac calcium channels: high-affinity binding to the inactivated state. *Proc Natl Acad Sci USA* 1984; 81: 6388-92.
- 12 Xu C, Sun MZ, Li YR, Yang BF, Wang LJ, Li JM. Inhibitory effect of tetrahydropalmatine on calcium current in isolated cardiomyocyte of guinea pig. *Acta Pharmacol Sin* 1996; 17: 329-31.

程 岚², 冯义柏, 夏国谨², 姚伟星²
(¹同济医科大学协和医院心血管疾病研究所心血管内科; ²同济医科大学基础医学院药理教研室, 武汉市 430022, 中国)

关键词 四氢巴马汀; 膜片箝技术; 钙电流; 心肌; 培养的细胞; 膜电压

目的: 研究左旋四氢巴马汀(L-THP)对L-钙电流的作用。方法: 用全细胞膜片箝记录心室肌细胞的L-钙电流。结果: L-THP 1-100 $\mu\text{mol} \cdot \text{L}^{-1}$ 将 $I_{Ca^{2+}-max}$ 从 (999 ± 93) pA 分别减少到 (700 ± 111) pA, (582 ± 66) pA 和 (420 ± 112) pA ($n=6, P < 0.01$)。L-THP 对钙通道有紧张性阻滞作用及使用依赖性。L-THP 可将钙通道的恢复时间从 (94 ± 39) ms 延长到 (170 ± 42) ms ($n=6, P < 0.01$)。结论: L-THP 对L-钙通道有阻滞作用。

(责任编辑 朱倩蓉)

左旋四氢巴马汀对单个豚鼠心室肌细胞钙电流的阻滞作用

黄 恺¹, 戴闰柱, 李新华, 范 曲²,

本部邮购科学出版社书讯(工具书类)

书 名	作 者	出书年月	单价(元)
汉英综合科学技术词汇(第二版)	科学出版社名词室	1997	88.00
英汉化学化工词汇(第三版)	科学出版社名词室	1997	68.00
生物学词典(第一版)	山田常雄(日), 等编, 鄂永昌, 等译	1997	150.00
现代免疫学词典	F.S. 罗森, 等编著, 朱立平, 等译	1997	16.00
新英汉病毒学词典	田波	1997	30.00
英汉—汉英生物化学词汇	沈昭文, 等	1996	25.00
细菌名称(第二版)	蔡妙英, 等	1996	55.00
英汉生物学词汇(第二版)	科学出版社名词室	1997	88.00
英汉常用生物学词汇	任久长, 崔克明, 等	1997	41.00
英汉生物化学词典	冯宗明	1983 年版(1997 重印)	58.00
英汉分子生物学与生物工程词汇	金冬雁, 等	1994	26.60

欲购者请另加 15 % 邮费, 汇款邮寄本部, 注明所购书名, 款到寄书。

联系地址: 200031 上海市太原路 294 号 《中国药理学报》编辑部 李慧珍 收。电话/传真: 021-6474-2629。