

## Inhibitory effect of aminoguanidine on bleomycin-induced pulmonary toxicity in rat

R916 A

CHEN Xiao-Ling<sup>1</sup>, HUANG Shan-Sheng, LI Wen-Bin, WANG Dian-Hua, WANG Xin-Liang (Department of Pathophysiology, Institute of Basic Medicine, Hebei Medical University, Shijiazhuang 050017, China)

**KEY WORDS** bleomycin; toxicity; pulmonary fibrosis; guanidines

### ABSTRACT

**AIM:** To observe the inhibitory effect of aminoguanidine (AG) on the pulmonary toxicity induced by bleomycin A<sub>5</sub> (BLM-A<sub>5</sub>) in rat. **METHODS:** The contents of hydroxyproline, nitrite/nitrate (NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup>), and malondialdehyde (MDA), which reflect fibrosis, nitric oxide (NO) production, and hyperoxidative injury of lung, were investigated by colorimetry. Histologic and morphometric examination of lung was also carried out on histological sections stained with hematoxylin and eosin (HE). **RESULTS:** (1) The content of hydroxyproline in lung and MDA in out-going pulmonary blood (OPB) increased from 14 d to 30 d after intratracheal administration of BLM-A<sub>5</sub>. Collapsed alveoli and fibrotic areas were seen and a lot of fibroblasts appeared in the lung interstitium of the rats 30 d after BLM-A<sub>5</sub>. (2) The NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> content increased in supernatant of alveolar macrophage culture and in OPB. (3) The increment of hydroxyproline in lung and MDA in OPB induced by BLM-A<sub>5</sub> was alleviated by aminoguanidine (AG, 20 mg·kg<sup>-1</sup>·d<sup>-1</sup>, ip). AG also reduced the histologic and morphometric changes in lung interstitium. (4) The augment of NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> in OPB was blocked by AG. **CONCLUSION:** AG had inhibitory effect on pulmonary injury and fibrosis induced by BLM-A<sub>5</sub>, and the inhibitory effect was related to the decrease of NO production in lung.

### INTRODUCTION

Bleomycin-induced pulmonary toxicity involved pul-

monary inflammatory injury and fibrosis. Despite the high efficiency of bleomycin-A<sub>5</sub> (BLM-A<sub>5</sub>) as a chemotherapeutic agent against various carcinomas, the pulmonary toxicity, especially the potentially lethal and chronic fibrotic response of the lung, is a major dose-limiting side effect<sup>[1,2]</sup>. So far, there was not effective therapy for the pulmonary fibrosis.

It has been reported that nitric oxide (NO) was involved in inflammatory response<sup>[3]</sup> and fibrosis of lung<sup>[4,5]</sup>. Quinlan *et al*<sup>[6]</sup> reported that NO generation might be important in cell injury and inflammation by asbestos. NO is produced by a five-electron oxidation of L-arginine to L-citrulline by NO synthase (NOS). Two general classes of NOS have been identified: the calcium-dependent, constitutively expressed NOS (cNOS) and the calcium-independent, transcriptionally inducible NOS (iNOS)<sup>[7]</sup>. It is well-accepted that cNOS was involved in physiological regulations. On the other hand, inflammatory stimuli, such as cytokines and endotoxin, upregulated iNOS. Produced in excess by iNOS, NO may cause tissue injury.

Aminoguanidine (AG), containing the guanido-group of L-arginine linked to hydrazine, *in vitro*, displays 10 to 100 folds higher potency as inhibitor of iNOS than cNOS<sup>[8]</sup>. In endotoxemic rodents and dogs, AG suppresses activation of iNOS and peroxynitrite production in the lungs and decreases plasma levels of the stable NO metabolites, nitrites, and nitrates. AG also reduces lung edema, improves gas exchange, and increases survival by counteracting circulatory failure<sup>[9-12]</sup>. Cruz *et al*<sup>[13,14]</sup> found that AG significantly attenuated the N-nitroso-N-methylurethane (NNMU)-induced alveolar injury. According above, We hypothesized that AG might suppress the progression of pulmonary injury and fibrosis induced by BLM-A<sub>5</sub>. Unfortunately, these have rarely been reported.

The current study was to prove enhancement of NO during pulmonary injury and fibrosis induced by BLM-A<sub>5</sub>, and to investigate the effect of AG in the above processes.

<sup>1</sup> Correspondence to Dr CHEN Xiao-Ling. Pbn 86-311-604-4121, ext 5607. E-mail chen\_xiaoling123@sina.com.cn  
Received 2001-01-22 Accepted 2001-04-21

## MATERIALS AND METHODS

**Experimental animals and groups** Sprague-Dawley rats (♂,  $n = 70$ , weighing 180–200 g, Grade II, Certificate No 06057) were provided by the Experimental Animal Center of Hebei Province, China. For the duration of the experiment, the animals were fed with commercial rat food and water *ad libitum*. The rats were randomly allocated to two treatment parts: (1) Rats received instillation of BLM- $A_5$  (5 mg/kg, in 0.5 mL of sterile saline) via tracheotomy under anesthesia (ip 2 % pentobarbital sodium 2 mL/kg) and sacrificed on d 7, d 14, d 21, and d 30, respectively (BLM- $A_5$  7 d, BLM- $A_5$  14 d, BLM- $A_5$  21 d, BLM- $A_5$  30 d group); or received intratracheal injection of the same amount of 0.9 % sterile saline as control group (Control); (2) Rats were given the same amount of BLM- $A_5$  as part (1), followed by daily intraperitoneal (ip) injection of AG (20 mg·kg<sup>-1</sup>·d<sup>-1</sup>) or saline for 14 d or 21 d. The rats were sacrificed on d 14 or d 30 after BLM- $A_5$  (BLM- $A_5$  + AG 14 d, BLM- $A_5$  + AG 30 d, BLM- $A_5$  + saline 14 d, BLM- $A_5$  + saline 14 d group).

**Collection of samples** Alveolar macrophages (from bronchoalveolar fluid) were placed in 96-well plates (in RPMI 1640 medium, 20 % fetal bovine serum,  $5 \times 10^8$  cells/L, 0.1 mL/well) and incubated for 24 h at 37 °C with 5 % CO<sub>2</sub>. After incubation, the culture supernatants of alveolar macrophage were used for detecting NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup>.

Out-going pulmonary blood (OPB) was collected under anesthesia (ip 2 % pentobarbital sodium 2 mL/kg) for detecting the nitrite/nitrate (NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup>) and malondialdehyde (MDA). Lung was taken for evaluating collagen by determining hydroxyproline and for morphometric examination.

### Measurements of nitrite/nitrate and MDA

The content of nitrite/nitrate (NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup>) in culture supernatants of alveolar macrophage and in OPB were measured by using detecting kit (Nanjing Jiancheng Biochemical Institute, China). The content of MDA in OPB was measured by using MDA detecting kit (Nanjing Jiancheng Biochemical Institute, China).

**Assay of hydroxyproline in lung** Hydroxyproline was released from lung tissue homogenates by acid hydrolysis (105 °C, 18 h). The free hydroxyproline was then oxidized by chloramine T to produce a pyrrole-type compound. The addition of Ehrlich's reagent (*p*-dimethylaminobenzaldehyde dissolved in perchloric

acid) resulted in the formation of a chromophore with a wavelength maximum at 550 nm. The absorbances were read using water as reference and corrected for the reagent blank. Absorbance values were plotted against nanomoles of added hydroxyproline and the negative intercept on the x axis was found by linear regression analysis. The absolute value of this intercept was taken to represent the hydroxyproline content of the tissue<sup>[15]</sup>.

**Histological and morphometric examination of lungs** Lungs from 6 rats of BLM- $A_5$  14 d and BLM- $A_5$  + AG 14 d groups were fixed by 4 % formaldehyde. Sections stained with hematoxylin and eosin (HE) were examined by light microscopy for evidence of pulmonary injury and fibrosis.

**Drugs** Bleomycin- $A_5$  (BLM- $A_5$ ) was produced by Tianjin Taihe Pharmaceutical Co, China. Aminoguanidine hemisulfate salt (AG) was produced in Germany, L-4-hydroxyproline was from Sigma Chemical Co, USA. RPMI 1640 was from GIBCO, USA. All other chemicals were of analytical reagent.

**Statistics** Data were expressed as  $\bar{x} \pm s$  and compared by *t*-test.

## RESULTS

**The toxicity of BLM- $A_5$  on lung** The degree of pulmonary fibrosis and injury was estimated by testing hydroxyproline in lung, MDA in OPB and histologic and morphometric changes of lungs. The contents of hydroxyproline in lung and MDA in OPB in BLM- $A_5$  14 d, 21 d, and 30 d groups were higher than those in the control group (Tab 1). Collapsed alveoli and fibrotic areas were seen and a lot of fibroblasts appeared in the lung interstitium of the rats in BLM- $A_5$  30 d group (Fig 1).

Tab 1. Hydroxyproline in lung and MDA and NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> in out-going pulmonary blood after intratracheal administration of BLM- $A_5$  in rats.  $n = 6$  rats.  $\bar{x} \pm s$ . <sup>a</sup>*P* < 0.05, <sup>b</sup>*P* < 0.01 vs control.

Groups	Hydroxyproline /mg·g <sup>-1</sup>	MDA /μmol·L <sup>-1</sup>	NO <sub>2</sub> <sup>-</sup> /NO <sub>3</sub> <sup>-</sup> /μmol·L <sup>-1</sup>
Control	19 ± 6	8.5 ± 1.5	60 ± 20
BLM- $A_5$ 7 d	24 ± 10	9.1 ± 1.1	103 ± 19 <sup>a</sup>
BLM- $A_5$ 14 d	33 ± 8 <sup>b</sup>	11.7 ± 0.8 <sup>b</sup>	171 ± 32 <sup>c</sup>
BLM- $A_5$ 21 d	52 ± 6 <sup>c</sup>	11.9 ± 1.0 <sup>b</sup>	77 ± 22
BLM- $A_5$ 30 d	56 ± 10 <sup>c</sup>	12.8 ± 0.5 <sup>c</sup>	77 ± 31




Fig 1. Typical morphology of lung tissue was observed in SD rat d 30 after intratracheal injection of BLM-A<sub>5</sub>. HE stain. × 400.

**Change of NO metabolites in lung induced by BLM-A<sub>5</sub>** The NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> in OPB were significantly increased in BLM-A<sub>5</sub> 7 d and BLM-A<sub>5</sub> 14 d group, and returned in BLM-A<sub>5</sub> 21 d and 30 d groups (Tab 1). Compared with control, the contents of NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> in culture supernatants of alveolar macrophages were high in BLM-A<sub>5</sub> 14 d and BLM-A<sub>5</sub> 30 d groups. But the NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> in supernatants in BLM-A<sub>5</sub> 30 d group was lower than that in BLM-A<sub>5</sub> 14 d group (Tab 2).

Tab 2. The content of NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> in culture supernatants of alveolar macrophages after administration of BLM-A<sub>5</sub> in rats. n = 4 rats. x ± s. <sup>b</sup>P < 0.05, <sup>c</sup>P < 0.01 vs control at the same day. <sup>a</sup>P < 0.05 vs BLM-A<sub>5</sub> d 14.

Groups	NO <sub>2</sub> <sup>-</sup> /NO <sub>3</sub> <sup>-</sup> (μmol·L <sup>-1</sup> )	
	d 14	d 30
Control	242 ± 32	241 ± 29
BLM-A <sub>5</sub>	442 ± 38 <sup>c</sup>	333 ± 38 <sup>bc</sup>

**Effects of AG on the pulmonary injury and fibrosis induced by BLM-A<sub>5</sub>** In BLM-A<sub>5</sub> + AG 30 d group, both hydroxyproline in lung and MDA in OPB were lower than those in BLM-A<sub>5</sub> + saline 30 d group. (Tab 3). In lung interstitium of rats in BLM-A<sub>5</sub> + AG 30 d group, the morphometric changes of BLM-A<sub>5</sub> 30 d group reduced (Fig 2).

**Effects of AG on NO in OPB of rats after administration of BLM-A<sub>5</sub>** In BLM-A<sub>5</sub> + AG 14 d group, NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> in OPB was lower than that in BLM-A<sub>5</sub> + saline 14 d group (Tab 4). There was no significant difference of the content of NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> in OPB between BLM-A<sub>5</sub> + AG 30 d group and BLM-A<sub>5</sub> + saline 30

d group (Tab 4).

Tab 3. Effects of AG on contents of hydroxyproline in lung and MDA in out-going pulmonary blood of rats on d 30 after intratracheal administration of BLM-A<sub>5</sub>. n = 6 rats. x ± s. <sup>a</sup>P < 0.01 vs BLM-A<sub>5</sub> + saline.

Groups	Hydroxyproline/ mg·g <sup>-1</sup>	MDA/μmol·L <sup>-1</sup>
BLM-A <sub>5</sub> + Saline	57 ± 4	12.5 ± 1.9
BLM-A <sub>5</sub> + AG	44 ± 4 <sup>a</sup>	8.0 ± 1.2 <sup>c</sup>




Fig 2. Typical morphology of lung tissue was observed in SD rat d 30 after intratracheal injection of BLM-A<sub>5</sub>, followed by ip AG 20 mg·kg<sup>-1</sup>·d<sup>-1</sup> for 21 d. HE stain. × 400.

Tab 4. Effects of AG on contents of NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> in out-going pulmonary blood of rats on d 14 and d 30 after intratracheal administration of BLM-A<sub>5</sub>. n = 6 rats. x ± s. <sup>a</sup>P < 0.01 vs BLM-A<sub>5</sub> + saline.

Groups	NO <sub>2</sub> <sup>-</sup> /NO <sub>3</sub> <sup>-</sup> (μmol·L <sup>-1</sup> )	
	d 14	d 30
BLM-A <sub>5</sub> + Saline	176 ± 17	96 ± 20
BLM-A <sub>5</sub> + AG	93 ± 19 <sup>a</sup>	80 ± 8

## DISCUSSION

In the present study, a large amount of NO was generated in lung during the period of hyperoxidative injury induced by BLM-A<sub>5</sub>. AG could block the above process and pulmonary fibrosis, as shown by the decreases of MDA level in OPB, hydroxyproline level in lung and reduction of the morphometric changes in lung interstitium. It was well known that NO was an important inflammatory mediator. High concentrations of NO may result in

greater accessibility of inflammatory cells to sites of injury<sup>[16]</sup>. Moreover, peroxyinitrite formed by the reaction of NO<sup>•</sup> with superoxide (O<sub>2</sub><sup>•-</sup>) may cause oxidant injury *per se* or after formation of a hydroxyl-like radical<sup>[17]</sup>. Saleh *et al*<sup>[18]</sup> had already found the increased production of peroxyinitrite in the lungs of patients with idiopathic pulmonary fibrosis. Paredi *et al*<sup>[5]</sup> found that exhaled NO was increased in active fibrosing alveolitis. These results suggested that NO might be one of the important factors to cause pulmonary injury. The inhibitory effect of AG might be related to block NO formation in lung.

Rezende *et al*<sup>[19]</sup> reported that after intratracheal administration of bleomycin, aminoguanidine bicarbonate (50 mg·kg<sup>-1</sup>·d<sup>-1</sup> for 4 weeks) led to significant reductions of hydroxyproline in murine lung, and the collagen area in the axial and septal zones of lung sections stained with Sirius red. In our experiment, the apparently high level of NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> in OPB appeared from d 7 to d 21 after injection of BLM-A<sub>5</sub>, and 20 mg·kg<sup>-1</sup>·d<sup>-1</sup> AG for 14 d could abolished above process. In addition, 20 mg·kg<sup>-1</sup>·d<sup>-1</sup> AG for 21 d could partly block the progression of pulmonary fibrosis. These results suggested that 20 mg·kg<sup>-1</sup>·d<sup>-1</sup> AG was enough to eliminate the influence of NO on pulmonary toxicity induced by BLM-A<sub>5</sub>.

Animal experiments listed some methods to modify the incidence and severity of pulmonary toxicity induced by BLM-A<sub>5</sub>. Anti-transforming growth factor-β antibodies<sup>[20-22]</sup> and human recombinant soluble tumor necrosis factor receptor<sup>[23]</sup> provided potentialities for intervening in the fibrotic process in the lung. *In vivo*, angiotensin-converting enzyme inhibitor captopril or the caspase inhibitor Z-Val-Ala-Asp-fluoromethylketone was used to block apoptosis of alveolar epithelial cell and lung fibrosis in response to BLM<sup>[24]</sup>. Keratinocyte growth factor, a potent proliferation and differentiation factor for rat alveolar type II cells, could prevent BLM-induced lung fibrosis in rats<sup>[25]</sup>. Our study was another way to prevent pulmonary toxicity induced by BLM. The level of NO in lung of patients could be estimated by NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> in out-going pulmonary blood. So, it's easy to determine the time when AG should be administered. This work demonstrated AG produced an inhibitory effect on pulmonary toxicity induced by BLM-A<sub>5</sub>. The observation suggests that AG may be a hopeful chemical reagent to prevent and cure the pulmonary fibrosis in-

duced by BLM-A<sub>5</sub>.

## REFERENCES

- Theile A, Kemper P. Bleomycin-induced pulmonary fibrosis following chemotherapy of ovarian granulosa cell tumor. *Pneumologie* 1998; 52: 325-32.
- Bloor AJ, Seale JR, Marcus RE. Two cases of fatal bleomycin pneumonitis complicating the treatment of non-Hodgkin's lymphoma. *Clin Lab Haematol* 1998; 20: 119-21.
- Ialenti A, Ianaro A, Moncada S, Di Rosa M. Modulation of acute inflammation by endogenous nitric oxide. *Eur J Pharmacol* 1992; 211: 177-82.
- Yamamoto T, Katayama I, Nishioka K. Nitrite production in mouse 3T3 fibroblasts by bleomycin-stimulated peripheral blood mononuclear cell factor. *Clin Exp Rheumatol* 1999; 17: 343-6.
- Paredi P, Kharitonov SA, Loukides S, Pantelidis P, de Bois RM, Barnes PJ. Exhaled nitric oxide is increased in active fibrosing alveolitis. *Chest* 1999; 115: 1352-6.
- Quinlan TR, Berube KA, Hacker MP, Taatjes DJ, Timblin CR, Goldberg J, *et al*. Mechanism of asbestos-induced nitric oxide production by rat alveolar macrophages in inhalation and *in vitro* models. *Free Radic Biol Med* 1998; 24: 778-88.
- Mayer B, Hemmens B. Biosynthesis and action of nitric oxide in mammalian cells. *Trends Biochem Sci* 1997; 22: 477-81.
- Misko TP, Moore WM, Kasten TP, Nickols GA, Corbett JA, Tilton RG, *et al*. Selective inhibition of the inducible nitric oxide synthase by aminoguanidine. *Eur J Pharmacol* 1993; 233: 119-25.
- Numata M, Suzuki S, Miyazawa N, Miyashita A, Nagashima Y, Inoue S, *et al*. Inhibition of inducible nitric oxide synthase prevents LPS-induced acute lung injury in dogs. *J Immunol* 1998; 160: 3031-7.
- Mikawa K, Nishina K, Tamada M, Takao Y, Maekawa N, Obara H. Aminoguanidine attenuates endotoxin-induced acute lung injury in rabbits. *Crit Care Med* 1998; 26: 905-11.
- Wu CC, Chen SJ, Szabo C, Thiemeermann C, Vane JR. Aminoguanidine attenuates the delayed circulatory failure and improves survival in rodent models of endotoxic shock. *Br J Pharmacol* 1995; 114: 1666-72.
- Arkovitz MS, Wispe JR, Garcia VF, Szabo C. Selective inhibition of the inducible isoform of nitric oxide synthase prevents pulmonary transvascular flux during acute endotoxemia. *J Pediatr Surg* 1996; 31: 1009-15.
- Cruz WS, Moxley MA, Corbett JA, Longmore WJ. Inhibition of nitric oxide synthase attenuates NNMU-induced alveolar injury *in vivo*. *Am J Physiol* 1997; 273 (6 Pt 1): L1167-73.
- Cruz WS, Corbett JA, Longmore WJ, Moxley MA. Nitric oxide participates in early events associated with NNMU-induced acute lung injury in rats. *Am J Physiol* 1999; 276 (2 Pt 1): L263-8.

- 15 Jamall IS, Finelli VN, Que Hee SS. A simple method to determine nanogram levels of 4-hydroxyproline in biological tissues. *Anal Biochem* 1981; 112: 70-5.
- 16 Kubes P, Granger DN. Nitric oxide modulates microvascular permeability. *Am J Physiol* 1992; 263: H611-5.
- 17 Beckman JS, Beckman TW, Chen J, Marshall PA, Freeman BA. Apparent hydroxyl radical production by peroxynitrite: Implications for endothelial injury from nitric oxide and superoxide. *Proc Natl Acad Sci USA* 1990; 87: 1620-4.
- 18 Saleh D, Barnes PJ, Giaid A. Increased production of the potent oxidant peroxynitrite in the lungs of patients with idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 1997; 155: 1763-9.
- 19 de Rezende MC, Martinez JA, Capelozzi VL, Simoes MJ, Beppu OS. Protective effect of aminoguanidine in a murine model of pulmonary fibrosis induced by bleomycin. *Fundam Clin Pharmacol* 2000; 14: 561-7
- 20 Giri SN, Hyde DM, Hollinger MA. Effect of antibody to transforming growth factor- $\beta$  on bleomycin induced accumulation of lung collagen in mice. *Thorax* 1993; 48: 959-66.
- 21 Wang Q, Wang Y, Hyde DM, Gotwals PJ, Kotliansky VE, Ryan ST, *et al.* Reduction of bleomycin induced lung fibrosis by transforming growth factor beta soluble receptor in hamsters. *Thorax* 1999; 54: 805-12.
- 22 Li J, He B, Weng B. The therapeutic effect of TGF-beta monoclonal antibody to bleomycin-induced pulmonary fibrosis in rats. *Chin J Tuberc Respir Dis* 1997; 20: 347-9.
- 23 Piguet PF, Vesin C. Treatment by human recombinant soluble TNF receptor of pulmonary fibrosis induced by bleomycin or silica in mice. *Eur Respir J* 1994; 7: 515-8.
- 24 Wang R, Ibarra-Sunga O, Verlinski L, Pick R, Uhal BD. Abrogation of bleomycin-induced epithelial apoptosis and lung fibrosis by captopril or by a caspase inhibitor. *Am J Physiol Lung Cell Mol Physiol* 2000; 279: L143-51.
- 25 Sugahara K, Lyama K, Kuroda MJ, Sano K. Double intratracheal instillation of keratinocyte growth factor prevents bleomycin-induced lung fibrosis in rats. *J Pathol* 1998; 186: 90-8.

## 氨基胍对博莱霉素诱发大鼠肺毒性的抑制作用

陈晓玲<sup>1</sup>, 黄善生, 李文斌, 王殿华, 王新良  
(河北医科大学基础医学研究所病生研究室, 石家庄 050017, 中国)

**关键词** 博莱霉素; 毒性; 肺纤维化; 胍类

**目的:** 观察氨基胍(AG)对博莱霉素 A<sub>5</sub>(BLMA<sub>5</sub>)诱发大鼠肺毒性作用的影响。 **方法:** BLM-A<sub>5</sub> (5 mg·kg<sup>-1</sup>)气管滴注诱发大鼠肺损伤和肺纤维化形成; 在气管滴注 BLM-A<sub>5</sub> 的同时及之后 21 d 腹腔注射 AG (20 mg·kg<sup>-1</sup>·d<sup>-1</sup>, ip)。 化学比色法测定肺组织羟脯氨酸含量、肺泡巨噬细胞培养上清液中 NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> 含量、出肺血中 NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> 含量; HE 染色组织切片上观察组织形态学变化。 **结果:** (1) 滴注 BLM-A<sub>5</sub> 后 d 14 至 30, 肺组织羟脯氨酸含量和出肺血中的 MDA 含量逐渐升高; 注 BLMA<sub>5</sub> 后 d 30, 肺间质可见萎陷的和纤维化的肺泡, 并出现大量的成纤维细胞。 (2) 滴注 BLM-A<sub>5</sub> 后 d 7 至 14, 出肺血 NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> 含量升高; 肺泡灌流液中巨噬细胞培养上清液中 NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> 含量升高。 (3) AG 明显抑制出肺血 MDA 含量和肺羟脯氨酸含量的升高; AG 还减轻肺间质组织形态学变化。 (4) AG 明显抑制出肺血 NO<sub>2</sub><sup>-</sup>/NO<sub>3</sub><sup>-</sup> 含量的升高。 **结论:** AG 对 BLM-A<sub>5</sub> 诱发的大鼠肺损伤和纤维化有抑制作用, 此作用与其抑制肺内 NO 大量生成有关。

(责任编辑 吕 静)